

## PHY-852 QUANTUM MECHANICS II

### Homework 4, 30 points

February 6 - 13, 2002

#### Motion in a central field.

Reading: *Merzbacher*, Chapter 11, sections 3 - 5; Chapter 12.

1. /8/ a. Find the first order differential equation for the spherical function  $Y_{il}(\theta, \phi)$  with the maximum projection  $m = l$ , solve this equation and normalize the solution. Explain the corresponding physical picture.  
b. Using the lowering operator construct the function  $Y_{l,l-1}(\theta, \phi)$ .
2. /4/ Consider a particle of mass  $m$  in a spherically symmetric potential well with the potential equal to  $U(r) = -U_0, U_0 > 0$ , for  $r < R$  and  $U(r) = 0$  for  $r > R$ . Find the interval of values  $U_0$  where the well supports two bound states with orbital momentum  $l = 0$ .
3. /7/ a. An object of mass  $M$  confines a loosely bound light particle of mass  $m$  with binding energy  $\epsilon$ . The particle is suddenly released from the system by an external perturbation, and the distribution of probabilities  $W(\mathbf{p})$  of different values of the particle momentum  $\mathbf{p}$  is measured. Assuming that the particle in the bound state spends most of the time outside the well, obtain a simple analytical expression for  $W(\mathbf{p})$ .  
b. In many experiments, instead of the full three-dimensional momentum distribution  $W(\mathbf{p})$ , only the "longitudinal" distribution  $W_x(p_x)$  of the momentum component  $p_x$  along a specific axis  $x$  is measured. Using the result a, predict the function  $W_x(p_x)$ .  
c. Assume that the neutron binding energy in a "halo" nucleus which consists of 3 protons and 8 neutrons is only 0.3 MeV. The experiment measuring  $W_x(p_x)$  shows a bell-shaped curve with the full width at half maximum equal to  $\Delta p_x = 44 \text{ MeV}/c$  where the convenient units for the momentum are used,  $c$  is the speed of light. Does this result agree with the predictions of the point b?
4. /5/ An electron is in the ground state of the tritium atom. The tritium nucleus  ${}^3\text{H}$  (one proton + two neutrons) suddenly undergoes beta-decay into the helium nucleus  ${}^3\text{He}$  (two protons and one neutron); the new electron and antineutrino created in the beta-decay carry away the electric charge and energy. Find the probability for the original atomic electron to remain in the ground state of the helium atom.
5. /6/ Consider the discrete spectrum of states of a particle in an attractive central field. Let  $E_{min}(l)$  be the lowest energy level with the orbital momentum  $l$ . Prove that

$$E_{min}(l) < E_{min}(l') \quad \text{if} \quad l < l'. \quad (1)$$