

PHY-852 QUANTUM MECHANICS II

Homework 6, 30 points

February 20 - 27, 2002

Elastic scattering and Born approximation.

Reading: *Merzbacher*, Chapters 13 and 20.

1. a. /3/ *Merzbacher*, Exercise 13.7.
- b. /5/ Establish the relation between your result and the Bragg condition

$$n\lambda = 2d \sin \frac{\theta}{2} \quad (1)$$

for the scattering angles θ that correspond to the interference maxima in the particle scattering by a system of parallel crystallic planes (n is an integer and λ is the wave length of the particle).

2. a. /3/ *Merzbacher*, Exercise 13.9.
 - b. /3/ *Merzbacher*, Exercise 13.10.
 - c. /4/ For the total cross section in problems a and b consider the limiting cases of low and high energies, $ka \ll 1$ and $ka \gg 1$, respectively (in the *Exercise 13.9* a is the radius of the well).
3. Consider the standard expression for the elastic scattering amplitude for a particle of mass m in the potential $U(\mathbf{r})$,

$$f(\mathbf{k}', \mathbf{k}) = -\frac{m}{2\pi\hbar^2} \int d^3r e^{-i(\mathbf{k}' \cdot \mathbf{r})} U(\mathbf{r}) \psi_{\mathbf{k}}(\mathbf{r}), \quad \mathbf{k}' = k \frac{\mathbf{r}}{r}, \quad (2)$$

where the wave function $\psi_{\mathbf{k}}(\mathbf{r})$ has at large distances r the asymptotic behavior

$$\psi_{\mathbf{k}}(\mathbf{r}) \approx e^{i(\mathbf{k} \cdot \mathbf{r})} + f(\mathbf{k}', \mathbf{k}) \frac{e^{ikr}}{r}. \quad (3)$$

- a. /4/ For the potential that at large distances falls off as $U(r) \propto r^{-s}$, find which values of s allow the asymptotic solution in the form (2).
- b. /5/ For a potential which becomes negligibly small at distances $r \sim R$, find at what distances the asymptotic representation (3) becomes valid.
- c. /3/ For an arbitrary potential $U(r)$ with central symmetry, show that if the Born approximation is valid, the product $E\sigma(E)$, where $\sigma(E)$ is the total scattering cross section at energy E , is a monotonously growing function of E .