

NAME.....

**PHY-852: QUANTUM MECHANICS I**

**Quiz 1**

*January 28, 2002*

**PROBLEM.** An angular part of the wave function of a particle is

$$\psi = A \sin^2 \theta. \tag{1}$$

Find the possible values of  $l$  and  $m$  in this state and their probabilities.

**SOLUTION.**

The absence of the  $\phi$ -dependence shows that only  $m = 0$  is possible. Because of parity, we can have only even  $l$ , and since this is a polynomial of the second order in  $\cos \theta$ , the allowed values are  $l = 0$  and  $l = 2$ . Therefore we have a superposition of  $Y_{00}$  and  $Y_{20}$ , or  $P_0 = 1$  and  $P_2 = (3/2) \cos^2 \theta - (1/2)$ ,

$$\psi = A[1 - \cos^2 \theta] = A \cdot \frac{2}{3}(P_0 - P_2) = \frac{2}{3}A\sqrt{4\pi} \left( Y_{00} - \frac{1}{\sqrt{5}}Y_{20} \right). \quad (2)$$

From the weights of the relative components of the orthonormalized functions  $Y_{00}$  and  $Y_{20}$  we find the probabilities  $w_2$  of  $l = 2$  and  $w_0$  of  $l = 0$ :

$$w_2 = \frac{1}{6}, \quad w_0 = \frac{5}{6}. \quad (3)$$