

NAME.....

**PHY-852: QUANTUM MECHANICS II**

**Quiz 4**

*April 2002*

**PROBLEM.** Electrons in an atom are in the stationary state with total angular momentum  $\mathbf{J}$ . A nucleus of the atom has its own angular momentum  $\mathbf{I}$ . The hyperfine interaction between the magnetic moments of the electrons and the nucleus is proportional to the scalar product of the angular momenta,

$$V_{h.f.s.} = A(\mathbf{J} \cdot \mathbf{I}). \quad (1)$$

a. Which of the following quantities are conserved by this interaction: components of the vectors  $\mathbf{J}, \mathbf{I}, \mathbf{J} \pm \mathbf{I}$ , their squares  $\mathbf{J}^2, \mathbf{I}^2, (\mathbf{J} \pm \mathbf{I})^2$ ?

b. For given absolute values of  $J$  and  $I$ , what splitting of original atomic levels is produced by this interaction (how many split levels and what is the energy spacing between them)?

**SOLUTION.** a. The total angular momentum of the atom

$$\mathbf{F} = \mathbf{J} + \mathbf{I} \quad (2)$$

is the only vector whose components are constants of motion. But all squares are conserved.

b. The split levels carry the possible values of  $F$  from  $|J - I|$  to  $J + I$ . The number of split levels is  $2J_{<} + 1$ , where  $J_{<}$  is the smaller of  $J$  and  $I$ . The shift of the level with the total momentum  $F$  is

$$\delta E(F) = \frac{A}{2}[F(F + 1) - J(J + 1) - I(I + 1)]. \quad (3)$$

The spacing between the adjacent split levels,

$$\delta E(F) - \delta E(F - 1) = \frac{A}{2}[F(F + 1) - (F - 1)F] = AF. \quad (4)$$

is linear in  $F$ .