

Many-Electron Atoms

Thornton and Rex, Ch. 8

In principle, can now solve Sch. Eqn for any atom.

In practice, -> Complicated!

Goal--

To explain properties of elements from principles of quantum theory
(without exact solutions)

- Elements distinguished by nuclear charge Z (= number of electrons)
- To first approx., each electron moves in electric field of nucleus + remaining electrons:

electron state : (n, ℓ, m_ℓ, m_s)

↗ shell ↖ subshell

{ n labels energy, but no simple formula.
 ℓ subshells no longer completely degenerate. }

- Principles for filling electron states:
 - 1) Always fill lowest energy state first.
 - 2) No two electrons can have same quantum numbers (n, ℓ, m_ℓ, m_s) .

Pauli Exclusion Principle

No two electrons can occupy the same quantum state.

Building up atomic structure of atoms

	<u>n</u>	<u>l</u>	<u>m_l</u>	<u>m_s</u>
Hydrogen	1	0	0	$\pm 1/2$

Helium	1	0	0	+1/2
	1	0	0	-1/2

Helium has a closed shell.

For Lithium, now add $n=2$ electron,
but $l=0$ or $l=1$?

Smaller l always has lower energy.

Lithium	1	0	0	+1/2
	1	0	0	-1/2
	2	0	0	$\pm 1/2$

	<u>n</u>	<u>l</u>	<u>m_l</u>	<u>m_s</u>
Hydrogen	1	0	0	+1/2
Helium	1	0	0	-1/2
Lithium	2	0	0	+1/2
Beryllium	2	0	0	-1/2
Boron	2	1	-1	+1/2
Carbon	2	1	0	+1/2
Nitrogen	2	1	+1	+1/2
Oxygen	2	1	-1	-1/2
Flourine	2	1	0	-1/2
Neon	2	1	+1	-1/2
Sodium	3	0	0	+1/2
Magnesium	3	0	0	-1/2
Aluminum	3	1	-1	+1/2
Silicon	3	1	0	+1/2
Phosporus	3	1	+1	+1/2
Sulfur	3	1	-1	-1/2
Chlorine	3	1	0	-1/2
Argon	3	1	+1	-1/2
Potassium	4	0	0	+1/2

(Last Electron Added)



Chemical properties of elements

Electrons in outermost, largest n orbits are most weakly bound. They determine the chemical properties of the elements. Elements with similar electron structure have similar properties.

- Inert or Noble Gases

Closed p subshell (s for He).

He ($1s^2$), Ne ($2s^22p^6$), Ar ($3s^23p^6$)

- Alkalis

Have single electron outside closed shell.

Li ($2s^1$), Na ($3s^1$), K ($4s^1$)

- Halogens

Are one electron short of a closed shell.

F ($2s^22p^5$), Cl ($3s^23p^5$)

Total Angular Momentum

Consider a 1-electron atom (or with just 1 electron outside closed shell).

It has Orbital Angular momentum \vec{L} and Spin Angular momentum \vec{S} .

These can be combined to give Total Angular momentum $\vec{J} = \vec{L} + \vec{S}$.

J is quantized with

$$J = \sqrt{j(j+1)} \hbar$$

and

$$J_z = m_j \hbar$$

where $j = \ell \pm s$

or $j = \ell \pm 1/2$ (since $s = 1/2$)

j will be half-integral ($1/2, 3/2, 5/2, \dots$)
 m_j will also be half-integral, ranging from $-j$ to j .

Example: $\ell=1, s=1/2$

$$m_\ell = (1, 0, -1) \quad m_s = (-1/2, +1/2)$$

$$3 \cdot 2 = 6 \text{ states}$$

Can combine into

$$j = 3/2 = 1 + 1/2$$

$$m_j = (-3/2, -1/2, +1/2, +3/2) \quad (4 \text{ states})$$


or

$$j = 1/2 = 1 - 1/2$$

$$m_j = (-1/2, +1/2) \quad (2 \text{ states})$$

$$\text{Total number of } j\text{-states is } 6 = 4 + 2.$$

Spectroscopic notation



Examples: $2S_{1/2}$ $3P_{3/2}$ etc.

Spin-Orbit Coupling

- Recall, coupling of spin to a magnetic field shifts the energy ($V_B = -\vec{\mu}_s \cdot \vec{B}$).
- Motion of electron produces an "internal" magnetic field.

So there is an additional contribution to the energy:

$$V_{SL} = -\vec{\mu}_s \cdot \vec{B}_{\text{int}}$$

Proportional to $-S$ Proportional to L


$$V_{SL} \propto \vec{S} \cdot \vec{L}$$

This is the Spin-Orbit Coupling:

$$V_{SL} = \mu \vec{S} \cdot \vec{L}$$


Now states with definite energy do not have unique L and S quantum numbers (m_ℓ, m_s). We must use J quantum numbers (j, m_j).

States with $j = \ell - 1/2$ have slightly less energy than states with $j = \ell + 1/2$.



(States with different m_j are still degenerate for each j .)

Selection Rules



Allowed transitions:

- lifetimes $\tau \sim 10^{-9}$ sec

$$\Delta n = \text{anything}, \quad \Delta \ell = \pm 1,$$

$$\Delta j = 0, \pm 1, \quad \Delta m_j = 0, \pm 1$$

Forbidden transitions:

- lifetimes much longer

$$\text{Ex. } 2s \rightarrow 1s, \tau \sim 1/7 \text{ sec}$$

Many-Electron Atoms

A careful analysis involving \vec{L} and \vec{S} in multi-electron atoms is very complicated.

Hund's Rules

(Empirical rules for filling a subshell, while minimizing the energy)

- 1) The total Spin should be maximized (without violating Pauli Exclusion Principle).
- 2) Without violating Rule 1, the Orbital Angular momentum should also be maximized.






Handwaving explanation:

Electrons repel each other, so we want them as far from each other as possible.

- 1) If spins of two electrons are aligned (for maximum \vec{S}), then Pauli Exclusion Principle says they must have different \vec{L} orbits. They will tend to be farther apart.
- 2) If the \vec{L} orbits are aligned (although with different magnitudes), then the electrons will travel around the nucleus in the same direction, so they don't pass each other as often.


Example:

A d subshell ($l=2$) can contain 10 electrons.



Example:

A d subshell ($l=2$) can contain 10 electrons.



Put first 5 electrons all with spin in same direction.

First 2 electrons have $m_l = -2$ and -1 , etc.

Many-Electron Atoms

For many-electron atoms there is now orbit-orbit and spin-spin interactions, in addition to spin-orbit interactions.

Consider simplest case of 2 electrons with \vec{L}_1, \vec{S}_1 and \vec{L}_2, \vec{S}_2 .

Only "good" quantum number is associated with total angular momentum

$$\vec{J} = \vec{L}_1 + \vec{L}_2 + \vec{S}_1 + \vec{S}_2 .$$

(By "good", I mean states with definite energy have definite j and m_j .)

How can we describe atom to best understand energy levels?

LS, or Russell-Saunders, Coupling

For most atoms the spin-orbit coupling is relatively weak. Then it makes sense to add the angular momentum in steps:

$$\text{First, } \vec{L} = \vec{L}_1 + \vec{L}_2$$
$$\vec{S} = \vec{S}_1 + \vec{S}_2$$

$$\text{Then } \vec{J} = \vec{L} + \vec{S}$$

For 2 electrons the Total Spin Quantum Number S is = 0 (spins anti-parallel)
or = 1 (spins parallel).

The Total Orbital Angular Momentum Quantum Number L is an integer in the range between $|l_1 - l_2|$ and $|l_1 + l_2|$.

The Total Angular Momentum Quantum Number J is an integer in the range between $|L - S|$ and $|L + S|$.

Note that for $S=0$, there is 1 value of J , given by $J=L$. This state is called a Singlet.

For $S=1$, there are 3 values of J , given by $J=L-1$, $J=L$, $J=L+1$. These states are called a Triplet.

In general, the multiplicity of the states is given by $(2S+1)$.

The Spectroscopic notation is

$$n^{(2S+1)}L_J$$

Example:

2 electrons, one in 4p, other in 4d.

I.e., $n=4$, $l_1=1$, $s_1=1/2$
 $l_2=2$, $s_2=1/2$

Possible values of S :

$$S=0 \quad \text{or} \quad S=1$$

Possible values of L :


$$L=1, 2, \text{ or } 3$$

Possible values of J :


$$\text{for singlet } (S=0): \quad J=L$$

$$\begin{aligned} \text{for triplet } (S=1): \quad & J=L-1 \\ & \text{or } J=L \\ & \text{or } J=L+1 \end{aligned}$$

Use Hund's rules to order the energies.



Use Hund's rules to order the energies.



Example:

Helium $1s^2$

$$l_1 = 0, s_1 = 1/2$$

$$l_2 = 0, s_2 = 1/2$$

Possible values of $S=0,1$

Possible values of $L=0$

Possible values of $J=0,1$

States: 1S_0, ~~3S_1~~

not allowed by Pauli Exclusion
(requires both electrons all same QN's)

If one electron is excited to $2s$, so the state is $1s2s$, then

both $^1S_0, ^3S_1$ are allowed.

jj Coupling

For high- Z elements the spin-orbit coupling is large for each electron.
Now add the angular momentum:

$$\text{First, } \vec{J}_1 = \vec{L}_1 + \vec{S}_1$$
$$\vec{J}_2 = \vec{L}_2 + \vec{S}_2$$

$$\text{Then } \vec{J} = \vec{J}_1 + \vec{J}_2$$

Anomalous Zeeman Effect

Recall, energy shift in external magnetic field:

$$V_B = -\vec{\mu} \cdot \vec{B}$$

The magnetic moment gets both orbital and spin contributions:

$$\vec{\mu} = \vec{\mu}_L + \vec{\mu}_S = \frac{-e}{2m} [\vec{L} + 2 \vec{S}]$$

If $S=0$, this is simple. It is just the Normal Zeeman effect. Energy levels split according to m_ℓ values:

$$V_B = m_\ell \mu_B B$$

But. most atoms are not "Normal".

If both S and L are nonzero, the spin-orbit coupling requires us to use J -states. Projecting \vec{L} onto \vec{J} gives

$$V_B = \frac{e}{2m} g \vec{J} \cdot \vec{B}$$
$$= \mu_B g m_J B$$

where the projection factor (called the Landé g factor) is

$$g = 1 + \frac{J(J+1) + S(S+1) - L(L+1)}{2J(J+1)}$$

This is the Anomalous Zeeman Effect.