Homework Set 3

Exercises; due Friday 9/18

- E5. Hand in a <u>computer generated</u> graph of $\sin \theta$ versus θ .
- E6. Write a short essay (2 paragraphs) on Synthetic Aperature Radar. The $1^{\rm st}$ paragraph should explain what it is. The $2^{\rm nd}$ paragraph should show an example an actual picture from the internet and explain how the image was obtained.

Problems; due Monday 9/21

- P11. A soap film with air on both sides has index of refraction n = 1.36. A region of the film appears red (w.l. = 633 nm) when white light is reflected from normal incidence. What is the thickness of the soap film in that region?
- P12. One flat glass plate is horizontal; a second plate rests on top of the first plate at an angle α . (Index of refraction of the glass = 1.5.) A parallel beam of monochromatic light (w.l. = 550 nm) is incident normally on the top plate. Calculate the angle α if the fringe separation is 0.25 cm.
- P13. As an example of thin film interference, consider light reflecting from an oil slick floating on water. Assume the thickness of the oil film is 500 molecular layers = 500×0.3 nm = 150 nm.
- (a) Assume n = 1.4 for the oil. Plot, accurately, a graph of the angle for the primary bright fringe versus the wavelength of light. [Use Mathematica or some other computer program to make the graph.]
- (b) For incident white light, what color would you see at grazing incidence?
- P14. Consider two-slit interference with these parameters:
- a = separation of the two slits = 0.02 mm; D = 2 m = distance from the slits to the observation screen; λ = wavelength = 550 nm.
- (a) Plot the intensity I(x) as a function of position x on the screen. (x = distance from the center to the point.)
- (b) Calculate the full width of the central bright fringe, i.e., between the two positions of minimum intensity.
- P15. Newton's rings. Suppose the diameter of the 5th bright ring is 5 mm.
- (a) Calculate the diameter of the 10th bright ring.
- (b) Calculate the distance (along a radial line) from the $10^{\rm th}$ bright ring to the $20^{\rm th}$ bright ring.
- P16. Newton's rings. <u>Use computer graphics</u>. Plot the bright rings --- a series of concentric rings with the correct radii --- for these parameters: R = 1 m, $\lambda = 550 \text{ nm}$.