Waves, the Wave Equation, and Phase Velocity

What is a wave?

Forward [f(x-vt)] and backward [f(x+vt)]propagating waves

The one-dimensional wave equation

Wavelength, frequency, period, etc.

Phase velocity Complex numbers

Plane waves and laser beams Boundary conditions

Div, grad, curl, etc., and the 3D Wave equation



Source: Trebino, Georgia Tech

What is a wave?

A wave is anything that moves.

To displace any function f(x) to the right, just change its argument from x to x-a, where a is a positive number.

If we let a = v t, where v is positive and t is time, then the displacement will increase with time.

So f(x - v t) represents a rightward, or forward, propagating wave.

Similarly, f(x + v t) represents a leftward, or backward, propagating wave.

v will be the velocity of the wave.





The one-dimensional wave equation

The one-dimensional wave equation for scalar (i.e., non-vector) functions, *f*:

$$\frac{\partial^2 f}{\partial x^2} - \frac{1}{\mathbf{v}^2} \frac{\partial^2 f}{\partial t^2} = 0$$

where v will be the velocity of the wave.

The wave equation has the simple solution:

$$f(x,t) = f(x \pm vt)$$

where f(u) can be any twice-differentiable function.

Proof that $f(x \pm vt)$ **solves the wave equation**

Write $f(x \neq vt)$ as f(u), where $u = x \neq vt$. So $\frac{\partial u}{\partial x} = 1$ and $\frac{\partial u}{\partial t} = \pm v$

Now, use the chain rule:
$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial x}$$
 $\frac{\partial f}{\partial t} = \frac{\partial f}{\partial u} \frac{\partial u}{\partial t}$

So
$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial u} \implies \frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial u^2}$$
 and $\frac{\partial f}{\partial t} = \pm v \frac{\partial f}{\partial u} \implies \frac{\partial^2 f}{\partial t^2} = v^2 \frac{\partial^2 f}{\partial u^2}$

Substituting into the wave equation:

$$\frac{\partial^2 f}{\partial x^2} - \frac{1}{v^2} \frac{\partial^2 f}{\partial t^2} = \frac{\partial^2 f}{\partial u^2} - \frac{1}{v^2} \left\{ v^2 \frac{\partial^2 f}{\partial u^2} \right\} = 0$$

The 1D wave equation for light waves

$$\frac{\partial^2 E}{\partial x^2} - \mu \varepsilon \frac{\partial^2 E}{\partial t^2} = 0$$

where *E* is the light electric field

We'll use cosine- and sine-wave solutions:

$$E(x,t) = B \cos[k(x \pm vt)] + C \sin[k(x \pm vt)]$$

$$\downarrow$$

$$kx \pm (kv)t$$

$$\downarrow$$

$$E(x,t) = B \cos(kx \pm \omega t) + C \sin(kx \pm \omega t)$$

or

where:

$$\frac{\omega}{k} = v = \frac{1}{\sqrt{\mu\varepsilon}}$$

The speed of light in vacuum, usually called "c", is 3×10^{10} cm/s.

$$E(x,t) = A \cos[(kx - \omega t) - \theta]$$

Use the trigonometric identity:

$$\cos(z-y) = \cos(z)\cos(y) + \sin(z)\sin(y)$$

where $z = kx - \omega t$ and $y = \theta$ to obtain:

$$E(x,t) = A \cos(kx - \omega t) \cos(\theta) + A \sin(kx - \omega t) \sin(\theta)$$

which is the same result as before,

$$E(x,t) = B\cos(kx - \omega t) + C\sin(kx - \omega t)$$

For simplicity, we'll just use the forward-propagating wave.

as long as:

$$A\cos(\theta) = B$$
 and $A\sin(\theta) = C$

Definitions: Amplitude and Absolute phase

$$E(x,t) = A \cos[(k x - \omega t) - \theta]$$

A = Amplitude

 θ = Absolute phase (or initial phase)



Definitions

Spatial quantities:



Temporal quantities:

The Phase Velocity

How fast is the wave traveling?

Velocity is a reference distance divided by a reference time.



The phase velocity is the wavelength / period: v = λ / τ

Since $v = 1/\tau$:

$$v = \lambda v$$

In terms of the k-vector, $k = 2\pi / \lambda$, and the angular frequency, $\omega = 2\pi / \tau$, this is:

$$v = \omega / k$$

The Phase of a Wave

The phase is everything inside the cosine.

$$E(x,t) = A \cos(\varphi)$$
, where $\varphi = k x - \omega t - \theta$

 $\varphi = \varphi(x, y, z, t)$ and is not a constant, like θ !

In terms of the phase,

$$\omega = -\partial \varphi / \partial t$$

$$k = \partial \varphi / \partial x$$

And

$$\mathbf{v} = \frac{-\partial \varphi / \partial t}{\partial \varphi / \partial x}$$

This formula is useful when the wave is really complicated.

Tacoma Narrows Bridge

Animation: http://www.youtube.com/watch?v=3mclp9QmCGs

1. The animation shows the Tacoma Narrows Bridge shortly before its collapse. What is its frequency?

- A .1 Hz
- B .25 Hz
- C .50 Hz
- D 1 Hz

2. The distance between the bridge towers (nodes) was about 860 meters and there was also a midway node. What was the wavelength of the standing torsional wave?

- A 1720 m
- B 860 m
- C 430 m
- D There is no way to tell.

3. What is the amplitude?

- A 0.4 m
- B 4 m
- C 8 m
- D 16 m

Complex numbers

Consider a point, P = (x,y), on a 2D Cartesian grid.



Let the x-coordinate be the real part and the y-coordinate the imaginary part of a complex number.

So, instead of using an ordered pair, (x,y), we write:

P = x + i y= $A \cos(\varphi) + i A \sin(\varphi)$

where $i = (-1)^{1/2}$

Euler's Formula

$$\exp(i\varphi) = \cos(\varphi) + i\sin(\varphi)$$

so the point, $P = A \cos(\varphi) + i A \sin(\varphi)$, can be written:

 $P = A \exp(i\varphi)$

where

A = Amplitude

 φ = Phase

Proof of Euler's Formula $\exp(i\varphi) = \cos(\varphi) + i\sin(\varphi)$

Use Taylor Series:
$$f(x) = f(0) + \frac{x}{1!}f'(0) + \frac{x^2}{2!}f''(0) + \frac{x^3}{3!}f'''(0) + \dots$$

$$\exp(x) = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$
$$\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} + \dots$$
$$\sin(x) = \frac{x}{1!} - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} + \dots$$

If we substitute $x = i\varphi$ into exp(x), then:

$$\exp(i\varphi) = 1 + \frac{i\varphi}{1!} - \frac{\varphi^2}{2!} - \frac{i\varphi^3}{3!} + \frac{\varphi^4}{4!} + \dots$$
$$= \left[1 - \frac{\varphi^2}{2!} + \frac{\varphi^4}{4!} + \dots\right] + i\left[\frac{\varphi}{1!} - \frac{\varphi^3}{3!} + \dots\right]$$
$$= \cos(\varphi) + i\sin(\varphi)$$

Complex number theorems

```
If \exp(i\varphi) = \cos(\varphi) + i\sin(\varphi)
             \exp(i\pi) = -1
             \exp(i\pi/2) = i
             \exp(-i\varphi) = \cos(\varphi) - i\sin(\varphi)
            \cos(\varphi) = \frac{1}{2} \left[ \exp(i\varphi) + \exp(-i\varphi) \right]
             \sin(\varphi) = \frac{1}{2i} \left[ \exp(i\varphi) - \exp(-i\varphi) \right]
             A_1 \exp(i\varphi_1) \times A_2 \exp(i\varphi_2) = A_1 A_2 \exp[i(\varphi_1 + \varphi_2)]
             A_1 \exp(i\varphi_1) / A_2 \exp(i\varphi_2) = A_1 / A_2 \exp[i(\varphi_1 - \varphi_2)]
```

More complex number theorems

Any complex number, *z*, can be written:

$$z = \operatorname{Re}\{z\} + i \operatorname{Im}\{z\}$$

So

and

Re{
$$z$$
 } = 1/2 ($z + z^*$)

Im{
$$z$$
 } = $1/2i$ ($z-z^*$)

where z^* is the complex conjugate of $z (i \rightarrow -i)$

The "magnitude," |z|, of a complex number is:

 $|z|^2 = z z^* = \operatorname{Re}\{z\}^2 + \operatorname{Im}\{z\}^2$

To convert z into polar form, $A \exp(i\varphi)$:

$$A^2 = \operatorname{Re}\{z\}^2 + \operatorname{Im}\{z\}^2$$

 $\tan(\varphi) = \operatorname{Im}\{z\} / \operatorname{Re}\{z\}$



We can also differentiate exp(ikx) as if the argument were real.

$$\frac{d}{dx}\exp(ikx) = ik\exp(ikx)$$

Proof:

$$\frac{d}{dx} \left[\cos(kx) + i\sin(kx) \right] = -k\sin(kx) + ik\cos(kx)$$

$$= ik \left[-\frac{1}{i}\sin(kx) + \cos(kx) \right]$$

But -1/i = i, so: $= ik [i \sin(kx) + \cos(kx)]$

Waves using complex numbers

The electric field of a light wave can be written:

$$E(x,t) = A \cos(kx - \omega t - \theta)$$

Since $\exp(i\varphi) = \cos(\varphi) + i \sin(\varphi)$, E(x,t) can also be written:

or

$$E(x,t) = \operatorname{Re} \left\{ A \exp[i(kx - \omega t - \theta)] \right\}$$

$$E(x,t) = \frac{1}{2} A \exp[i(kx - \omega t - \theta)] + c.c.$$
We often
write these
expressions
without the
¹/₂, Re, or
+c.c.

where "+ c.c." means "plus the complex conjugate of everything before the plus sign."

Waves using complex amplitudes

We can let the amplitude be complex:

$$E(x,t) = A \exp[i(kx - \omega t - \theta)]$$
$$E(x,t) = \{A \exp(-i\theta)\} \{\exp[i(kx - \omega t)]\}$$

where we've separated the constant stuff from the rapidly changing stuff.

The resulting "complex amplitude" is:

So:

$$\underline{E}_0 = A \exp(-i\theta) \quad \leftarrow \text{(note the " ~ ")}$$

$$E(x, t) = E \exp(i(kx - \omega t)) \quad \text{As } x$$

 $E(x,t) = E_0 \exp i(kx - \omega t)$ As written, this entire field is complex!

How do you know if E_0 is real or complex?

Sometimes people use the "~", but not always. So always assume it's complex. Adding waves of the same frequency, but different initial phase, yields a wave of the same frequency.

This isn't so obvious using trigonometric functions, but it's easy with complex exponentials:

$$\begin{split} \tilde{E}_{tot}(x,t) &= \tilde{E}_1 \exp i(kx - \omega t) + \tilde{E}_2 \exp i(kx - \omega t) + \tilde{E}_3 \exp i(kx - \omega t) \\ &= (\tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3) \exp i(kx - \omega t) \end{split}$$

where all initial phases are lumped into E_1 , E_2 , and E_3 .



$E_0 \exp[i(kx - \omega t)]$ is called a plane wave.

A plane wave's contours of maximum field, called **wave-fronts** or **phase-fronts**, are planes. They extend over all space.

Wave-fronts are helpful for drawing pictures of interfering waves.



A wave's wavefronts sweep along at the speed of light.

A plane wave's wave-fronts are equally spaced, a wavelength apart.

They're perpendicular to the propagation direction.

Usually, we just draw lines; it's easier.

Localized waves in space: beams

A plane wave has flat wave-fronts throughout all space. It also has infinite energy. It doesn't exist in reality.

Real waves are more localized. We can approximate a realistic wave as a plane wave vs. z times a Gaussian in x and y:

$$E(x, y, z, t) = E_0 \exp\left[-\frac{x^2 + y^2}{w^2}\right] \exp[i(kz - \omega t)]$$





Localized waves in time: pulses

If we can localize the beam in space by multiplying by a Gaussian in x and y, we can also localize it in time by multiplying by a Gaussian in time.



$$\underbrace{E}_{\tilde{\omega}}(x, y, z, t) = \underbrace{E}_{0} \exp\left[-\frac{t^{2}}{\tau^{2}}\right] \exp\left[-\frac{x^{2} + y^{2}}{w^{2}}\right] \exp[i(kz - \omega t)]$$

This is the equation for a laser pulse.

Longitudinal vs. Transverse waves



Space has 3 dimensions, of which 2 are transverse to the propagation direction, so there are 2 transverse waves in addition to the potential longitudinal one.

The direction of the wave's variations is called its polarization.

Vector fields

Light is a 3D vector field.

A 3D vector field $\vec{f}(\vec{r})$ assigns a 3D vector (i.e., an arrow having both direction and length) to each point in 3D space.



Wind patterns: 2D vector field

A light wave has both electric and magnetic 3D vector fields:



And it can propagate in any direction.

Div, Grad, Curl, and all that

Types of 3D vector derivatives:

The Del operator:
$$\vec{\nabla} \equiv \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}\right)$$

The Gradient of a scalar function f:

$$\vec{\nabla}f \equiv \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}\right)$$

div grad an curl informal text and on vector all calculus that third edition h.m.schey

Div, Grad, Curl, and All That: An Informal Text on Vector Calculus , by Schey

> If you want to know more about vector calculus, read this book!

The gradient points in the direction of steepest ascent.

Div, Grad, Curl, and all that

The **Divergence** of a vector function:

$$\vec{\nabla} \cdot \vec{f} \equiv \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} + \frac{\partial f_z}{\partial z}$$

The **Divergence** is nonzero if there are sources or sinks.

A 2D source with a large divergence:



Div, Grad, Curl, and more all that

The Laplacian of a scalar function :

$$\nabla^2 f \equiv \nabla \cdot \nabla f = \nabla \cdot \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}\right)$$
$$= \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

The Laplacian of a vector function is the same, but for each component of *f*:

$$\nabla^{2}\vec{f} = \left(\frac{\partial^{2}f_{x}}{\partial x^{2}} + \frac{\partial^{2}f_{x}}{\partial y^{2}} + \frac{\partial^{2}f_{x}}{\partial z^{2}}\right), \quad \frac{\partial^{2}f_{y}}{\partial x^{2}} + \frac{\partial^{2}f_{y}}{\partial y^{2}} + \frac{\partial^{2}f_{y}}{\partial z^{2}}, \quad \frac{\partial^{2}f_{z}}{\partial x^{2}} + \frac{\partial^{2}f_{z}}{\partial y^{2}} + \frac{\partial^{2}f_{z}}{\partial z^{2}}\right)$$

The Laplacian tells us the curvature of a vector function.

The 3D wave equation for the electric field and its solution

A light wave can propagate in any direction in space. So we must allow the space derivative to be 3D:

$$\vec{\nabla}^2 E - \mu \varepsilon \frac{\partial^2 E}{\partial t^2} = 0$$

or

$$\frac{\partial^2 E}{\partial x^2} + \frac{\partial^2 E}{\partial y^2} + \frac{\partial^2 E}{\partial z^2} - \mu \varepsilon \frac{\partial^2 E}{\partial t^2} = 0$$

whose solution is:

$$E(x, y, z, t) = \operatorname{Re}\left\{ \tilde{E}_0 \exp[i(\vec{k} \cdot \vec{r} - \omega t)] \right\}$$

where
$$\vec{k} \equiv (k_x, k_y, k_z)$$
 $\vec{r} \equiv (x, y, z)$

and $\vec{k} \cdot \vec{r} \equiv k_x x + k_y y + k_z z$

$$k^2 \equiv k_x^2 + k_y^2 + k_z^2$$

The 3D wave equation for a light-wave electric field is actually a vector equation.

And a light-wave electric field can point in any direction in space:

$$\vec{\nabla}^2 \vec{E} - \mu \varepsilon \frac{\partial^2 \vec{E}}{\partial t^2} = 0$$

Note the arrow over the *E*.

whose solution is:
$$\vec{E}(x, y, z, t) = \operatorname{Re}\left\{\vec{E}_{0} \exp[i(\vec{k} \cdot \vec{r} - \omega t)]\right\}$$

where:
 $\vec{E}_{0} = (E_{0x}, E_{0y}, E_{0z})$

Vector Waves

We must now allow the field *E* and its complex field amplitude E_0 to be vectors:

$$\vec{E}(\vec{r},t) = \operatorname{Re}\left\{\vec{E}_{0} \exp\left[i\left(\vec{k}\cdot\vec{r}-\omega t\right)\right]\right\}$$

The complex vector amplitude has six numbers that must be specified to completely determine it!



Boundary Conditions

Often, a wave is constrained by external factors, which we call **Boundary Conditions**.

For example, a guitar string is attached at both ends.

In this case, only certain wavelengths/frequencies are possible.

Here the wavelengths can be:

 λ_1 , $\lambda_1/2$, $\lambda_1/3$, $\lambda_1/4$, etc.



