

Hertzprung-Russell Diagram, Flux, Luminosity, Magnitude—10 Oct

- Outline
 - Review of 7 Oct
 - Thermal radiation
 - Wien's Law
 - Stefan-Boltzmann Law
 - How to measure temperature of stars. AJ Cannon's method of classifying spectra.
 - Hertzprung-Russell diagram
 - There are 3 types of stars: main-sequence or dwarfs, giants, white dwarfs
 - New material
 - H-R diagram of star clusters
 - Flux, luminosity, magnitude, absolute magnitude

Thermal Radiation

- Spectrum (intensity vs. wavelength) of thermal radiation.

- Hotter objects are brighter at all wavelengths.

- Wien's Law

$$\lambda_{\text{peak}} T = 2.9 \text{ mm K}$$

- Wavelength changes inversely with temperature

- For the sun, $T=5700\text{K}$.

$$\lambda_{\text{peak}} = \frac{2.9 \text{ mm K}}{5700 \text{ K}} = 0.0005 \text{ mm} = 500 \text{ nm}$$

- For a person, $T=273+37=310\text{K}$.


$$\lambda_{\text{peak}} = \frac{2.9 \text{ mm K}}{310 \text{ K}} = 0.01 \text{ mm} = 10 \mu\text{m}$$

- Stefan-Boltzmann Law

- Energy emitted per second depends on area A (or radius R for a sphere) and temperature T


$$L = AT^4$$

$$L = R^2 T^4$$



Luminosity & Flux (apparent brightness) of Stars

- Luminosity is amount of energy per second (Watt) produced by the star.
 - Intrinsic to the star.
 - With constants suppressed,
 $L = R^2 T^4$
- Flux is energy per second received by a detector on earth (Watt/m²).
 - Depends on distance of star
 $F = L/D^2$
 - At greater distances from star, light is spread over larger area. Flux is lower.




Is it flux or luminosity?

- The color of my cat is a property of my cat. It does not change with distance. I see the same color whether my cat is 1' or 10' from me.
1. S1: The flux of a star does not change with distance.
S2: The luminosity does not change with distance.
 - A. TT
 - B. TF
 - C. FT
 - D. FF
 2. ___ is the quantity that I measure directly.
 - A. Flux
 - B. Luminosity

Applying the hot-plate model to stars

- Flux=Luminosity/distance²
- Luminosity=R²T⁴
- It is around 1900.
 - The distances to some stars are known. They were measured by the method of _____. That means the luminosities are known.
 - To figure out the hot-plate model of a star, a measurement of _____ is needed.
- Annie Jump Cannon figure out a way to measure it.

Annie Jump Cannon: Classify stars by spectra



- Classification is very efficient: Draper catalog has 250,000 stars.
- Spectral class was later found to be related to temperature.
O be a fine girl kiss me.

- Prof. Pickering's Team in 1913, from Barbara L. Welther, 1982, Isis 73, 94.
- AJC
 - BA, Wellesley, 1884
 - Pickering's assistant, 1896
 - Henry Draper catalog of stars, 1918-1924
 - Astronomer 1938



Hertzsprung-Russell diagram


- H-R Diagram is plot of temperature & luminosity
 - Hotter stars are on left.
 - More luminous stars are on the top.
 - Stars exist only with certain combinations of luminosity and temperature.



Ejnar Hertzsprung
1873-1967 (Danish)




Henry Norris Russell
1877-1957 (American)



Astrophysics and twentieth-century astronomy to 1950, O Gingerich, ed., Cambridge, 1984

Hertzsprung-Russell diagram


- H-R Diagram is plot of temperature & luminosity
 - Stefan-Boltzmann Law: $L = AT^4$
1. A star gets hotter and its size does not change. In the H-R diagram, it moves
- up & left
 - up & right
 - up-down
 - left-right
 - not at all



Astrophysics and twentieth-century astronomy to 1950, O Gingerich, ed., Cambridge, 1984

Hertzsprung-Russell diagram


- H-R Diagram is plot of temperature & luminosity
 - Stefan-Boltzmann Law: $L = AT^4$
1. Can two stars of the same spectral class have different luminosities?
- No. No such cases exist on the H-R diagram.
 - Yes, temperatures differ
 - Yes, sizes differ
 - Yes, both size & temperatures differ.



Astrophysics and twentieth-century astronomy to 1950, O Gingerich, ed., Cambridge, 1984

Hertzsprung-Russell diagram

- H-R Diagram is plot of temperature & luminosity
- Stefan-Boltzmann Law:
 $L = AT^4$
- H-R diagram reveals stars cannot have any combination of size and temperature. There are three types of stars.
- Dwarfs have differing temperatures and approximately the same size.
 - Dwarfs are most common.
 - Also called main-sequence stars.
- Giants are large.
- White dwarfs are small.



Astrophysics and twentieth-century astronomy to 1950,
O Gingerich, ed., Cambridge, 1984