

Physics 842 – Fall 2012

Classical Electrodynamics II

Notation differences between Landau & Lifshitz and PHY842 lectures & homeworks

Physical quantity Landau & Lifshitz PHY842 lectures and homeworks

Chapters 1 and 2:

electric charge	e	q
dielectric susceptibility	κ	χ _e

Chapter 3, section 26 has some confusing definitions:

chemical potential	ζ	μ
internal chemical potential	ζ ₀	μ _{int}
energy flux density	q	j_ε
heat current	q - φj	j_q = j_ε - μj_n

Landau & Lifshitz redefine φ on page 97 as φ + ζ₀/e to include the chemical potential. The new φ introduced by L&L is equal to μ/q in my notation. The modern way to view this is to say that the total chemical potential is a sum of an “internal” term and an “external” term: μ = μ_{int} + μ_{ext} = μ_{int} + qφ, where φ is the usual electrostatic potential. (The external term could also include the effect of gravity or any other external force.)

thermopower	α	S
-------------	---	---

Chapter 4:

total current density	ρv	j or j_{total}
contributions to current	ρv = c∇× M + j	j_{total} = c∇×M + j_{free}
surface current density	g	J

(I do not like the L & L use of ρv for total current density, since “bound currents” include electric spins, which cannot be thought of as a charge density ρ moving at velocity v.)