


Quiz 1


1. C&J page 6, Check Your Understanding #3 –
Select an equation that DOES NOT have the same units on the two sides of the equal sign.

2. Two vectors, \vec{A} and \vec{B} , are shown in the diagram. What are the signs of the components of the vector, $\vec{C} = \vec{A} + \vec{B}$?



- a) C_x is +, and C_y is + b) C_x is +, and C_y is –
c) C_x is –, and C_y is + d) C_x is –, and C_y is –

3. A car travels 30° South of West for 240 miles. How far has it gone to the South?




- a) 120 miles, b) 66 miles c) 100 miles d) 80 miles e) 150 miles

Quiz 1

1. C&J page 6, Check Your Understanding #3 –
Select an equation that DOES NOT have the same units
on the two sides of the equal sign.


d) or e)

2. Two vectors, \vec{A} and \vec{B} , are shown
in the diagram. What are the signs
of the components of the vector,
 $\vec{C} = \vec{A} + \vec{B}$?



- a) C_x is +, and C_y is + b) C_x is +, and C_y is –
c) C_x is –, and C_y is + d) C_x is –, and C_y is –

3. A car travels 30° South of West for 240
miles. How far has it gone to the South?



- a) 120 miles, b) 66 miles c) 100 miles d) 80 miles e) 150 miles

Chapter 2

Kinematics in One Dimension

Kinematics deals with the concepts that are needed to describe motion.

Dynamics deals with the effect that forces have on motion.

Together, kinematics and dynamics form the branch of physics known as ***Mechanics***.

2.0 Definitions

Speed v : is a positive scalar
Velocity \vec{v} : is a vector.

} Instantaneous - at a time t .

Magnitude of the velocity vector is the speed, v .

Direction - for projectiles (2 or 3D), direction is an angle.

- for motion along a line (1D), direction is a sign.


$$\vec{v} = +v \quad \text{or} \quad \vec{v} = -v$$

2.0 Definitions, continued


In Chapter 2: All motion is along a line. It could be horizontal (cars, boats, humans), or vertical (acceleration due to gravity) **MUST** decide which direction is going to be POSITIVE

1D Examples:


Sliding block




OR



motion of block
thrown upward



OR



2.0 Definitions, continued

Moving: How can one tell if an object is moving at time, t ?

Look a little bitty time (ϵ) earlier, $t' = t - \epsilon$,

or look a little bitty time (ϵ) later, $t' = t + \epsilon$

and see if the object is at the same place.


If object is at the same place, it is not moving (stationary).

If object is NOT at the same place --- it is MOVING.

If an object is thrown upward, at the highest point
 $v = 0$, instantaneously, but the object IS MOVING!

Turning around is motion, it is moving.


2.1 Displacement



\vec{x}_0 = initial position (at $t = 0$) \vec{x} = final position

$$\Delta\vec{x} = \vec{x} - \vec{x}_0 = \text{displacement}$$


2.1 Displacement



$$\Delta\vec{x} = \vec{x} - \vec{x}_o = 7.0 \text{ m} - 2.0 \text{ m} = 5.0 \text{ m}$$

(Note: $\vec{x}_o + \Delta\vec{x} = \vec{x}$ initial position + displacement = final position)

2.1 Displacement



$$\Delta\vec{x} = \vec{x} - \vec{x}_o = 2.0 \text{ m} - 7.0 \text{ m} = -5.0 \text{ m}$$

2.1 Displacement



$$\Delta\vec{x} = 7.0 \text{ m}$$

(displacement)

$$\Delta\vec{x} = \vec{x} - \vec{x}_o = 5.0 \text{ m} - (-2.0) \text{ m} = 7.0 \text{ m}$$

2.2 *Speed and Velocity*

Average speed is the distance traveled divided by the time required to cover the distance.

$$\text{Average speed} = \frac{\text{Distance}}{\text{Elapsed time}}$$

SI units for speed: **meters per second** (m/s)

2.2 *Speed and Velocity*

Example 1 Distance Run by a Jogger

How far does a jogger run in 1.5 hours (5400 s) if his average speed is 2.22 m/s?

$$\text{Average speed} = \frac{\text{Distance}}{\text{Elapsed time}}$$

$$\begin{aligned} \text{Distance} &= (\text{Average speed})(\text{Elapsed time}) \\ &= (2.22 \text{ m/s})(5400 \text{ s}) = 12000 \text{ m} \end{aligned}$$

Clicker Question 2.1

A jogger travels 1500 m at an average speed of 2.00 m/s.
How long did it take to cover the distance ?

- a) 300 seconds
- b) 750 seconds
- c) 3000 seconds
- d) 1700 seconds
- e) 10 minutes

Clicker Question 2.1

A jogger travels 1500 m at an average speed of 2.00 m/s.
How long did it take to cover the distance ?

a) 300 seconds

b) 750 seconds

c) 3000 seconds

d) 1700 seconds

e) 10 minutes

Average speed

$$\bar{v} = \frac{d \text{ (distance)}}{t \text{ (elapsed time)}}$$

$$t = \frac{d}{\bar{v}} = \frac{1500 \text{ m}}{2.00 \text{ m/s}} = 750 \text{ s}$$

2.2 *Speed and Velocity*

Average velocity is the displacement divided by the elapsed time.

$$\text{Average velocity} = \frac{\text{Displacement}}{\text{Elapsed time}}$$

$$\vec{\mathbf{v}} = \frac{\vec{\mathbf{x}} - \vec{\mathbf{x}}_o}{t - t_o} = \frac{\Delta \vec{\mathbf{x}}}{\Delta t}$$



A vector divided by a scalar is another vector

Note: division by a 2D or 3D vector is not allowed

2.2 Speed and Velocity


Example 2 The World's Fastest Jet-Engine Car

Andy Green in the car *ThrustSSC* set a world record of 341.1 m/s in 1997. To establish such a record, the driver makes two runs through the course, one in each direction, to nullify wind effects. From the data, determine the average velocity for each run.




2.2 Speed and Velocity

$$\vec{v} = \frac{\Delta \vec{x}}{\Delta t} = \frac{+1609 \text{ m}}{4.740 \text{ s}} = +339.5 \text{ m/s}$$



$$\vec{v} = \frac{\Delta \vec{x}}{\Delta t} = \frac{-1609 \text{ m}}{4.695 \text{ s}} = -342.7 \text{ m/s}$$




2.2 *Speed and Velocity*

The ***instantaneous velocity*** indicates how fast the car moves and the direction of motion at each instant of time.


$$\vec{v} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{x}}{\Delta t}$$

2.3 Acceleration

The notion of *acceleration* emerges when a change in velocity is combined with the time during which the change occurs.




2.3 Acceleration



DEFINITION OF AVERAGE ACCELERATION

$$\vec{a} = \frac{\vec{v} - \vec{v}_o}{t - t_o} = \frac{\Delta \vec{v}}{\Delta t} \quad \left(\begin{array}{l} \text{rate that velocity} \\ \text{is changing in time} \end{array} \right)$$

2.3 Acceleration




Example 3 Acceleration and Increasing Velocity

Determine the average acceleration of the plane.

$$\vec{v}_o = 0 \text{ m/s} \quad \vec{v} = 260 \text{ km/h} \quad t_o = 0 \text{ s} \quad t = 29 \text{ s}$$

$$\bar{\mathbf{a}} = \frac{\vec{v} - \vec{v}_o}{t - t_o} = \frac{260 \text{ km/h} - 0 \text{ km/h}}{29 \text{ s} - 0 \text{ s}} = +9.0 \frac{\text{km/h}}{\text{s}}$$



2.3 Acceleration



2.3 Acceleration

Example 3 Acceleration and Decreasing Velocity

$$\bar{\mathbf{a}} = \frac{\vec{\mathbf{v}} - \vec{\mathbf{v}}_o}{t - t_o} = \frac{13 \text{ m/s} - 28 \text{ m/s}}{12 \text{ s} - 9 \text{ s}} = -5.0 \text{ m/s}^2$$




$$\bar{\mathbf{a}} = -5.0 \text{ m/s}^2$$



(b)

2.3 Acceleration



Acceleration was $\vec{a} = -5.0 \text{ m/s}^2$ throughout the motion

2.4 Equations of Kinematics for Constant Acceleration

$$\vec{\mathbf{v}} = \frac{\vec{\mathbf{x}} - \vec{\mathbf{x}}_o}{t - t_o}$$

$$\vec{\mathbf{a}} = \frac{\vec{\mathbf{v}} - \vec{\mathbf{v}}_o}{t - t_o}$$

For motion along a straight line

it is customary to **dispense** with the use of boldface symbols overdrawn with arrows for the displacement, velocity, and acceleration vectors. We will, however, continue to convey the directions with a plus or minus sign.

$$v = \frac{x - x_o}{t - t_o}$$

$$a = \frac{v - v_o}{t - t_o}$$

a, v and v_o are STILL vectors (values will have + or – for direction)

2.4 Equations of Kinematics for Constant Acceleration


From now on unless stated otherwise

Let the object be at the origin when the clock starts.

$$x_o = 0 \quad (t_o = 0)$$

Simplifies things a great deal

$$\bar{v} = \frac{x - x_o}{t - t_o} \quad \longrightarrow \quad \bar{v} = \frac{x}{t}$$



Note: average is
(initial + final)/2

$$x = \bar{v}t = \frac{1}{2}(v_o + v)t$$


2.4 Equations of Kinematics for Constant Acceleration

A constant acceleration can
be measured at any time


$$a = \frac{v - v_o}{t - t_o}$$



$$a = \frac{v - v_o}{t}$$



$$at = v - v_o$$



$$v = v_o + at$$


2.4 *Equations of Kinematics for Constant Acceleration*

Five kinematic variables:


1. displacement, x
2. acceleration (constant), a
3. final velocity (at time t), v
4. initial velocity, v_0
5. elapsed time, t

2.4 Equations of Kinematics for Constant Acceleration

$$v = v_o + at$$




$$x = \frac{1}{2} (v_o + v)t = \frac{1}{2} (v_o + v_o + at)t$$




$$x = v_o t + \frac{1}{2} at^2$$

2.4 Equations of Kinematics for Constant Acceleration



$$\begin{aligned}x &= v_0 t + \frac{1}{2} a t^2 \\&= (6.0 \text{ m/s})(8.0 \text{ s}) + \frac{1}{2} (2.0 \text{ m/s}^2)(8.0 \text{ s})^2 \\&= +110 \text{ m}\end{aligned}$$

2.4 Equations of Kinematics for Constant Acceleration



(b)

Example 6 Catapulting a Jet

Find its displacement.

$$v_o = 0 \text{ m/s}$$


$$x = ??$$

$$a = +31 \text{ m/s}^2$$


$$v = +62 \text{ m/s}$$

2.4 Equations of Kinematics for Constant Acceleration

$$a = \frac{v - v_o}{t} \quad \longrightarrow \quad t = \frac{v - v_o}{a}$$




$$x = \frac{1}{2} (v_o + v) t = \frac{1}{2} (v_o + v) \frac{(v - v_o)}{a}$$



$$x = \frac{v^2 - v_o^2}{2a}$$

2.4 Equations of Kinematics for Constant Acceleration



(b)

$$x = \frac{v^2 - v_o^2}{2a} = \frac{(62 \text{ m/s})^2 - (0 \text{ m/s})^2}{2(31 \text{ m/s}^2)} = +62 \text{ m}$$

2.4 *Equations of Kinematics for Constant Acceleration*

Equations of Kinematics for Constant Acceleration

$$v = v_o + at$$

$$x = \frac{1}{2} (v_o + v) t$$

$$v^2 = v_o^2 + 2ax$$

$$x = v_o t + \frac{1}{2} at^2$$

2.5 *Applications of the Equations of Kinematics*

Reasoning Strategy

1. Make a drawing.
2. Decide which directions are to be called positive (+) and negative (-).
3. Write down the values that are given for any of the five kinematic variables.
4. Verify that the information contains values for at least three of the five kinematic variables. Select the appropriate equation.
5. When the motion is divided into segments, remember that the final velocity of one segment is the initial velocity for the next.
6. Keep in mind that there may be two possible answers to a kinematics problem.

Clicker Question 2.2

A car accelerates from rest to 100 m/s in 400 meters.
What was the acceleration of the car in this motion?

Hint: use $v^2 = v_o^2 + 2ax$

- a) 12.5 m/s²
- b) 4.00 m/s²
- c) 0.25 m/s²
- d) 400 m/s²
- e) 25 m/s²

Clicker Question 2.2

A car accelerates from rest to 100 m/s in 400 meters.
What was the acceleration of the car in this motion?

Hint: use $v^2 = v_0^2 + 2ax$

a) 12.5 m/s²

b) 4.00 m/s²

c) 0.25 m/s²

d) 400 m/s²

e) 25 m/s²

$$v_0 = 0; \quad v = 100 \text{ m/s}; \quad x = 400 \text{ m}$$

$$a = \frac{v^2}{2x} = \frac{10,000 \text{ m}^2/\text{s}^2}{2(400 \text{ m})} = 12.5 \text{ m/s}^2$$


2.5 Applications of the Equations of Kinematics

Example 8 An Accelerating Spacecraft

A spacecraft is traveling with a velocity of +3250 m/s. Suddenly the **retrorockets** are fired, and the spacecraft begins to **slow down** with an acceleration whose **magnitude** is 10.0 m/s^2 . What is the velocity of the spacecraft when the displacement of the craft is +215 km, relative to the point where the retrorockets began firing?

x	a	v	v_o	t
+215000 m	-10.0 m/s ²	?	+3250 m/s	

2.5 Applications of the Equations of Kinematics



2.5 Applications of the Equations of Kinematics

x	a	v	v_o	t
+215000 m	-10.0 m/s ²	?	+3250 m/s	

$$v^2 = v_o^2 + 2ax \longrightarrow v = \sqrt{v_o^2 + 2ax}$$

$$\begin{aligned} v &= \pm \sqrt{(3250 \text{ m/s})^2 + 2(-10.0 \text{ m/s}^2)(215000 \text{ m})} \\ &= \pm 2500 \text{ m/s} \end{aligned}$$

2.6 *Freely Falling Bodies*

In the absence of air resistance, it is found that all bodies at the same location above the Earth fall vertically with the same acceleration. If the distance of the fall is small compared to the radius of the Earth, then the acceleration remains essentially constant throughout the descent.

This idealized motion is called *free-fall* and the acceleration of a freely falling body is called the *acceleration due to gravity*.

$$g = 9.80 \text{ m/s}^2 \quad \text{or} \quad 32.2 \text{ ft/s}^2$$

2.6 Freely Falling Bodies



Air-filled
tube
(a)



Evacuated
tube
(b)

*acceleration due
to gravity.*

$$g = 9.80 \text{ m/s}^2$$