

Quiz on Chapters 13-15

Final Exam, Thursday May 3, 8:00 – 10:00PM
ANH 1281 (Anthony Hall). Seat assignments TBD

RCPD students: Thursday May 3, 5:00 – 9:00PM, BPS
3239. Email will be sent.

Alternate Final Exam, Tuesday May 1, 10:00AM – noon,
BPS 3239; BY APPOINTMENT ONLY, and deadline has
past. Email will be sent.

Quiz 11

1. C&J p. 393 (top), Check Your Understanding #8: “Two bars are ...”

a) $Q_1 = \frac{1}{4} Q_2$ b) $Q_1 = \frac{1}{8} Q_2$ c) $Q_1 = 2Q_2$ d) $Q_1 = 4Q_2$ e) $Q_1 = Q_2$

Quiz 11

1. C&J p. 393 (top), Check Your Understanding #8: “Two bars are ...”

a) $Q_1 = \frac{1}{4}Q_2$ b) $Q_1 = \frac{1}{8}Q_2$ c) $Q_1 = 2Q_2$ d) $Q_1 = 4Q_2$ e) $Q_1 = Q_2$

2. C&J p. 419 (mid), Check Your Understanding #15: “The pressure of ...”

$$v = v_{rms} = \sqrt{3kT / m}$$

a) $\frac{v_f}{v_i} = 2$ b) $\frac{v_f}{v_i} = \sqrt{2}$ c) $\frac{v_f}{v_i} = \frac{1}{2}$ d) $\frac{v_f}{v_i} = \frac{1}{\sqrt{2}}$ e) $\frac{v_f}{v_i} = 1$

Quiz 11

1. C&J p. 393 (top), Check Your Understanding #8: “Two bars are ...”

a) $Q_1 = \frac{1}{4}Q_2$ b) $Q_1 = \frac{1}{8}Q_2$ c) $Q_1 = 2Q_2$ d) $Q_1 = 4Q_2$ e) $Q_1 = Q_2$

2. C&J p. 419 (mid), Check Your Understanding #15: “The pressure of ...”

$$v = v_{rms} = \sqrt{3kT / m}$$

a) $\frac{v_f}{v_i} = 2$ b) $\frac{v_f}{v_i} = \sqrt{2}$ c) $\frac{v_f}{v_i} = \frac{1}{2}$ d) $\frac{v_f}{v_i} = \frac{1}{\sqrt{2}}$ e) $\frac{v_f}{v_i} = 1$

3. What thickness of concrete, with a thermal conductivity of 2.2 J/(s•m•K) will conduct heat at the same rate as 0.25 m of air, which has a thermal conductivity of 0.0256 J/(s•m•K), if all other conditions are the same?

- a) 8.9 m
- b) 4.3 m
- c) 1.4 m
- d) 11 m
- e) 22 m

$$\frac{Q}{t} = \frac{kA\Delta T}{L}$$

$\frac{Q}{t}$: rate of heat flow

k : thermal conductivity

A : area

ΔT : temperature difference

L : thickness

Quiz 11

4. What mass of Carbon Dioxide (CO₂) has the same number of ATOMS as 0.100 kg of Oxygen (O₂). (atomic weight of C = 12u, of O = 16u).
- a) 0.032 kg
 - b) 0.12 kg
 - c) 0.062 kg
 - d) 0.044 kg
 - e) 0.096 kg

Quiz 11

4. What mass of Oxygen (O_2) has the same number of ATOMS as 0.088 kg of Carbon Dioxide (CO_2) . (atomic mass of C = 12 u, and of O = 16 u). .
- a) 0.032 kg
 - b) 0.12 kg
 - c) 0.062 kg
 - d) 0.044 kg
 - e) 0.096 kg
5. A cylinder with a moveable piston contains an ideal gas. The gas is subsequently expanded **adiabatically**. Which of the following choices correctly identifies the signs of (Q) the heat exchanged with the environment, (W) the work done, and (ΔU) the change in the internal energy?
- a) $Q = 0$, $W = -$, $\Delta U = -$
 - b) $Q = -$, $W = +$, $\Delta U = -$
 - c) $Q = 0$, $W = -$, $\Delta U = +$
 - d) $Q = 0$, $W = +$, $\Delta U = -$
 - e) $Q = +$, $W = -$, $\Delta U = 0$

Quiz 11

1. C&J p. 393 (top), Check Your Understanding #8: “Two bars are ...”

- a) $Q_1 = \frac{1}{4}Q_2$ b) $Q_1 = \frac{1}{8}Q_2$ **c) $Q_1 = 2Q_2$** d) $Q_1 = 4Q_2$ e) $Q_1 = Q_2$

$$Q_2 \propto k_2 A_2 = \left(\frac{1}{6}k_1\right)(3A_1) = \frac{1}{2}Q_1 \Rightarrow (Q_1 = 2Q_2)$$

1. C&J p. 419 (mid), Check Your Understanding #15: “The pressure of ...”

- a) $\frac{v_f}{v_i} = 2$ b) $\frac{v_f}{v_i} = \sqrt{2}$ c) $\frac{v_f}{v_i} = \frac{1}{2}$ **d) $\frac{v_f}{v_i} = \frac{1}{\sqrt{2}}$** e) $\frac{v_f}{v_i} = 1$

$$P_f = 2P_i, \quad V_f = \frac{1}{4}V_i$$

$$P_f V_f = \frac{1}{2}P_i V_i = nR\left(\frac{1}{2}T_i\right), \quad \underline{T_f = \frac{1}{2}T_i}$$

$$v_f = \sqrt{3kT_f / m}$$

$$= \frac{1}{\sqrt{2}} \sqrt{3kT_i / m} = \frac{1}{\sqrt{2}} v_i$$

2. What thickness of concrete, with a thermal conductivity of 2.2 J/(s•m•K) will conduct heat at the same rate as 0.25 m of air, which has a thermal conductivity of 0.0256 J/(s•m•K), if all other conditions are the same?

- a) 8.9 m
b) 4.3 m
c) 1.4 m
d) 11 m
e) 22 m

$$\frac{k_A A \Delta T}{L_A} = \frac{k_C A \Delta T}{L_C} \quad (\text{air and concrete heat rates are the same})$$

$$\frac{k_A}{L_A} = \frac{k_C}{L_C} \Rightarrow L_C = \frac{k_C}{k_A} L_A = \frac{2.2}{0.0256} (0.25\text{m}) = \underline{22\text{ m}}$$

Quiz 11

4. What mass of Oxygen (O₂) has the same number of ATOMS as 0.088 kg of Carbon Dioxide (CO₂) . (atomic mass of C = 12 u, and of O = 16 u).

a) 0.032 kg

b) 0.12 kg

c) 0.062 kg

d) 0.044 kg

e) 0.096 kg

$$n_{\text{O}_2} = 0.088 \text{ kg} / (0.044 \text{ kg/mole}) = 2.0 \text{ moles of CO}_2$$

$$N_{\text{atoms}} = (2)n_{\text{O}_2}N_A = (3)n_{\text{CO}_2}N_A \Rightarrow 3n_{\text{CO}_2} = 2n_{\text{O}_2}$$

$$n_{\text{O}_2} = \frac{3}{2}n_{\text{CO}_2} = 3.0 \text{ moles}$$

$$m_{\text{O}_2} = n_{\text{O}_2} (0.032 \text{ kg}) = 3.0(0.032 \text{ kg}) = \underline{0.096 \text{ kg}}$$

5. A cylinder with a moveable piston contains an ideal gas. The gas is subsequently expanded **adiabatically**. Which of the following choices correctly identifies the signs of (Q) the heat exchanged with the environment, (W) the work done, and (ΔU) the change in the internal energy?

a) $Q = 0$, $W = -$, $\Delta U = -$

b) $Q = -$, $W = +$, $\Delta U = -$

c) $Q = 0$, $W = -$, $\Delta U = +$

d) $Q = 0$, $W = +$, $\Delta U = -$

e) $Q = +$, $W = -$, $\Delta U = 0$

adiabatically $\Rightarrow Q = 0$

expanded $\Rightarrow W$ is positive

$\Delta U = -W \Rightarrow \Delta U$ is negative

Chapter 16

Waves and Sound

continued

Final Exam, Thursday May 3, 8:00 – 10:00PM
ANH 1281 (Anthony Hall). Seat assignments TBD

RCPD students: Thursday May 3, 5:00 – 9:00PM, BPS
3239. Email will be sent.

Alternate Final Exam, Tuesday May 1, 10:00AM – noon,
BPS 3239; BY APPOINTMENT ONLY, and deadline has
past. Email will be sent.

16.8 Decibels

The **decibel** (dB) is a measurement unit used when comparing two sound Intensities.


Human hearing mechanism responds to sound **intensity level** , logarithmically.

$$\beta = (10 \text{ dB}) \log\left(\frac{I}{I_o}\right)$$

Note that $\log(1) = 0$

dB (decibel)

$$I_o = 1.00 \times 10^{-12} \text{ W/m}^2$$



	Intensity I (W/m^2)	Intensity Level β (dB)
Threshold of hearing	1.0×10^{-12}	0
Rustling leaves	1.0×10^{-11}	10
Whisper	1.0×10^{-10}	20
Normal conversation (1 meter)	3.2×10^{-6}	65
Inside car in city traffic	1.0×10^{-4}	80
Car without muffler	1.0×10^{-2}	100
Live rock concert	1.0	120
Threshold of pain	10	130

16.8 Decibels


Example 9 Comparing Sound Intensities

Audio system 1 produces a sound intensity level of 90.0 dB, and system 2 produces an intensity level of 93.0 dB. Determine the ratio of intensities.

$$\beta = (10 \text{ dB}) \log\left(\frac{I}{I_o}\right)$$

$$\begin{aligned} 90 \text{ dB} &= (10 \text{ dB}) \log(I/I_o) \\ \log(I/I_o) &= 9; \\ I &= I_o \times 10^9 = (1 \times 10^{-12} \text{ W/m}^2) \times 10^9 \\ &= 1 \times 10^{-3} \text{ W/m}^2 \end{aligned}$$

$$\begin{aligned} 93 \text{ dB} &= (10 \text{ dB}) \log(I/I_o) \\ \log(I/I_o) &= 9.3; \\ I &= I_o \times 10^{9.3} = (1 \times 10^{-12} \text{ W/m}^2) \times 10^{9.3} \\ &= 1 \times 10^{-2.7} \text{ W/m}^2 = 1 \times 10^{-3} (10^{0.3}) \text{ W/m}^2 \\ &= 1 \times 10^{-3} (2) \text{ W/m}^2 = 2 \times 10^{-3} \text{ W/m}^2 \end{aligned}$$



$$93 \text{ dB} = 90 \text{ dB} + 3 \text{ dB}$$

Adding 3dB results in a factor of 2 increase in the intensity.

$$3 \text{ dB} = (10 \text{ dB}) \log(I_2/I_1)$$

$$0.3 = \log(I_2/I_1);$$

$$I_2 = 10^{0.3} I_1 = 2 I_1$$

Clicker Question 16.1

Software is used to amplify a digital sound file on a computer by 30 dB. By what factor has the intensity of the sound been increased as compared to the original sound file?

$$\beta = (10 \text{ dB}) \log \left(\frac{I}{I_o} \right)$$

- a) 1000
- b) 100
- c) 20
- d) 5
- e) 2

Clicker Question 16.1

Software is used to amplify a digital sound file on a computer by 30 dB. By what factor has the intensity of the sound been increased as compared to the original sound file?

a) 1000

b) 100

c) 20

d) 5


e) 2

$$\begin{aligned}\beta_2 &= \beta_1 + 30 \text{ dB} \\ (10 \text{ dB}) \log\left(\frac{I_2}{I_o}\right) &= (10 \text{ dB}) \log\left(\frac{I_1}{I_o}\right) + 30 \text{ dB} \\ \log I_2 &= \log I_1 + 3 \\ I_2 &= 10^{\log I_1 + 3} = 10^{\log I_1} \cdot 10^3 \\ &= \underline{10^3} I_1\end{aligned}$$


Take the dB increase and divide by 10.

The intensity increase factor is 10 to that power.

16.9 The Doppler Effect



(a)



(b)

The **Doppler effect** is the change in frequency or pitch of the sound detected by an observer because the sound source and the observer have different velocities with respect to the medium of sound propagation.

**SOURCE (s) MOVING AT v_s
TOWARD OBSERVER (o)**

$$f_o = f_s \left(\frac{1}{1 - v_s/v} \right)$$

**SOURCE (s) MOVING AT v_s
AWAY FROM OBSERVER (o)**


$$f_o = f_s \left(\frac{1}{1 + v_s/v} \right)$$

Chapter 17


The Principle of Linear Superposition and Interference Phenomena

17.1 The Principle of Linear Superposition

When the pulses merge, the Slinky assumes a shape that is the sum of the shapes of the individual pulses.



(a) Overlap begins




(b) Total overlap; the Slinky has twice the height of either pulse




(c) The receding pulses

17.1 The Principle of Linear Superposition

When the pulses merge, the Slinky assumes a shape that is the sum of the shapes of the individual pulses.



(a) Overlap begins



(b) Total overlap




(c) The receding pulses


17.1 The Principle of Linear Superposition

THE PRINCIPLE OF LINEAR SUPERPOSITION

When two or more waves are present simultaneously at the same place, the resultant disturbance is the sum of the disturbances from the individual waves.




(a) Overlap begins




(b) Total overlap; the Slinky has twice the height of either pulse



(c) The receding pulses



(a) Overlap begins




(b) Total overlap



(c) The receding pulses


17.2 Constructive and Destructive Interference of Sound Waves

When two waves always meet condensation-to-condensation and rarefaction-to-rarefaction, they are said to be **exactly in phase** and to exhibit **constructive interference**.




17.2 Constructive and Destructive Interference of Sound Waves

When two waves always meet condensation-to-rarefaction, they are said to be **exactly out of phase** and to exhibit **destructive interference**.




17.2 Constructive and Destructive Interference of Sound Waves



17.2 Constructive and Destructive Interference of Sound Waves

If the wave patterns do not shift relative to one another as time passes, the sources are said to be **coherent**.

For two wave sources vibrating in phase, a difference in path lengths that is zero or an integer number (1, 2, 3, . . .) of wavelengths leads to constructive interference; a difference in path lengths that is a half-integer number ($\frac{1}{2}$, $1 \frac{1}{2}$, $2 \frac{1}{2}$, . . .) of wavelengths leads to destructive interference.




17.2 Constructive and Destructive Interference of Sound Waves

Example 1 What Does a Listener Hear?

Two in-phase loudspeakers, A and B, are separated by 3.20 m. A listener is stationed at C, which is 2.40 m in front of speaker B.

Both speakers are playing identical 214-Hz tones, and the speed of sound is 343 m/s.

Does the listener hear a loud sound, or no sound?



Calculate the path length difference.

$$\sqrt{(3.20 \text{ m})^2 + (2.40 \text{ m})^2} - 2.40 \text{ m} = 1.60 \text{ m}$$

Calculate the wavelength.

$$\lambda = \frac{v}{f} = \frac{343 \text{ m/s}}{214 \text{ Hz}} = 1.60 \text{ m}$$

Because the path length difference is equal to an integer (1) number of wavelengths, there is constructive interference, which means there is a loud sound.





17.2 *Constructive and Destructive Interference of Sound Waves*

Conceptual Example 2 Out-Of-Phase Speakers



To make a speaker operate, two wires must be connected between the speaker and the amplifier. To ensure that the diaphragms of the two speakers vibrate in phase, it is necessary to make these connections in exactly the same way. If the wires for one speaker are not connected just as they are for the other, the diaphragms will vibrate out of phase. Suppose in the figures (next slide), the connections are made so that the speaker diaphragms vibrate out of phase, everything else remaining the same. In each case, what kind of interference would result in the overlap point?



17.2 Constructive and Destructive Interference of Sound Waves




17.3 Diffraction



The bending of a wave around an obstacle or the edges of an opening is called ***diffraction***.

17.3 Diffraction



single slit – first minimum $\sin \theta = \frac{\lambda}{D}$

Clicker Question 17.1

Sound with a wavelength of 2.8 m is directed at a doorway that is 3.2 m wide. Due to diffraction, the sound will be nearly zero intensity at what angle?

$$\sin \theta = \frac{\lambda}{D}$$

- a) 83°
- b) 72°
- c) 61°
- d) 50°
- e) 30°

Clicker Question 17.1

Sound with a wavelength of 2.8 m is directed at a doorway that is 3.2 m wide. Due to diffraction, the sound will be nearly zero intensity at what angle?

a) 83°

b) 72°

c) 30°


d) 52°

e) 61°

$$\sin \theta = \frac{\lambda}{D}$$

$$\theta = \sin^{-1} \left(\frac{\lambda}{D} \right) = \sin^{-1} \left(\frac{2.8}{3.2} \right) = \sin^{-1}(0.875)$$
$$= 61^\circ$$


17.3 Diffraction



Circular opening – first minimum



$$\sin \theta = 1.22 \frac{\lambda}{D}$$

17.4 Beats



Two overlapping waves with *slightly different frequencies* gives rise to the phenomena of beats.






17.4 Beats




The **beat frequency** is the **difference** between the two sound frequencies.

17.5 Transverse Standing Waves

Transverse standing wave patterns.



17.5 Transverse Standing Waves








In reflecting from the wall, a forward-traveling half-cycle becomes a backward-traveling half-cycle that is inverted.

Unless the timing is right, the newly formed and reflected cycles tend to offset one another.

Repeated reinforcement between newly created and reflected cycles causes a large amplitude standing wave to develop.



17.5 Transverse Standing Waves



String fixed at both ends

$$f_n = n \left(\frac{v}{2L} \right) \quad n = 1, 2, 3, 4, \dots$$

17.6 Longitudinal Standing Waves



Tube open at both ends

$$f_n = n \left(\frac{v}{2L} \right) \quad n = 1, 2, 3, 4, \dots$$


17.6 Longitudinal Standing Waves

Example 6 Playing a Flute



When all the holes are closed on one type of flute, the lowest note it can sound is middle C (261.6 Hz). If the speed of sound is 343 m/s, and the flute is assumed to be a cylinder open at both ends, determine the distance L .

$$f_n = n \left(\frac{v}{2L} \right) \quad n = 1, 2, 3, 4, \dots$$

$$L = \frac{nv}{2f_n} = \frac{1(343 \text{ m/s})}{2(261.6 \text{ Hz})} = 0.656 \text{ m}$$



17.6 Longitudinal Standing Waves



Tube open at one end

$$f_n = n \left(\frac{v}{4L} \right) \quad n = 1, 3, 5, \dots$$

17.7 Complex Sound Waves

