

your name(s) _____

Physics 852 Quiz #16 - Friday, March. 20th

Coulomb's Law

In Chapter 9 the electromagnetic field operator was written as,

$$\vec{A}(\vec{r}, t) = \sqrt{\frac{2\pi\hbar^2 c^2}{V}} \sum_{\vec{k}, s} \vec{\epsilon}_s(\vec{k}) \frac{1}{\sqrt{E(\vec{k})}} \left(e^{i\vec{k} \cdot \vec{r} - iE(\vec{k})t/\hbar} a_{\vec{k}, s} + e^{-i\vec{k} \cdot \vec{r} + iE(\vec{k})t/\hbar} a_{\vec{k}, s}^\dagger \right).$$

This ignores the zeroth component of \vec{A} because there are only two polarizations, and neither has a zeroth component. Although there are no physical, on-shell, phonons with polarizations either parallel to \vec{k} , or in the zeroth direction, one can write the field with all four polarizations,

$$A_\mu(\vec{r}, t) = \sqrt{\frac{2\pi\hbar^2 c^2}{V}} \sum_{\vec{k}} \frac{1}{\sqrt{E(\vec{k})}} \left(e^{ik \cdot x} a_{\mu, \vec{k}} + e^{-ik \cdot x} a_{\mu, \vec{k}}^\dagger \right).$$

There is no longer any need to write down polarization vectors, because the the creation/destruction operators now behave as four vectors. Note that x refers to the four-vector t, \vec{r} and $k \cdot x = Et/\hbar - \vec{k} \cdot \vec{r}$. The commutation rules are

$$[a_{\mu, \vec{k}}, a_{\nu, \vec{k}'}^\dagger] = -g_{\mu\nu} \delta_{\vec{k}\vec{k}'}.$$

For $\mu = 1, 2, 3$ these are the usual commutation rules because $-g_{ii} = 1$. However, the signs are reversed for the zeroth component, as if the probability of that state is negative (more on that later).

Your mission, should you choose to accept it, is to consider a static charge density,

$$j_0(\vec{r}) = \rho(\vec{r}) \neq 0, \vec{j}(\vec{r}) = 0,$$

and calculate the energy due to interaction with the quantum electromagnetic field in second order perturbation theory. Here, the charge density is fixed (or classical), and can be considered as a given function of x , i.e. it is not altered by the interaction with the electromagnetic field. The state $|0\rangle$ refers to a state with zero photons (the ground state), but it does have the static charge density. The state $|\mu, \vec{k}\rangle$ refers to a field with the same charge density but with a single photon of wave vector \vec{k} and polarization μ . Note that if the state $|\mu, \vec{k}\rangle$ is to have unit norm, then if $|\mu, \vec{k}\rangle = a_{\mu, \vec{k}}^\dagger |0\rangle$ the bra must be defined as $\langle\mu, \vec{k}| = -\langle 0| a_{\mu, \vec{k}}$. Equivalently, for $\mu = 0$, $a_{\mu=0}^\dagger = -(a_{\mu=0})^\dagger$.

The correction to the ground state energy in second order perturbation theory is

$$\begin{aligned} \delta E^{(2)} &= - \sum_{i \neq 0} \frac{|\langle\mu, \vec{k}| H_{\text{int}} |0\rangle|^2}{E_i}, \\ H_{\text{int}} &= \int d^3r j^\alpha(\vec{r}) A_\alpha(\vec{r}), \\ E(\vec{k}) &= \hbar c k. \end{aligned}$$

1. Using the definitions above, show that $\delta E^{(2)}$ can be written as

$$\delta E^{(2)} = \int d^3x_1 d^3x_2 \frac{\hbar^2 c^2}{(2\pi)^2} \int \frac{d^3k}{E(\vec{k})^2} \rho(\vec{x}_1) \rho(\vec{x}_2) e^{i\vec{k} \cdot (\vec{x}_1 - \vec{x}_2)}.$$

2. Perform the integral over \vec{k} to derive Coulomb's law,

$$\delta E^{(2)} = \int d^3x_1 d^3x_2 \frac{1}{2|\vec{x}_1 - \vec{x}_2|} \rho(\vec{x}_1) \rho(\vec{x}_2).$$

Note that the energy is positive (for any $\rho(\vec{x})$). However, from the expression for $\delta E^{(2)}$ one would expect that the second-order correction to the ground state energy should always be negative. The source of this paradox comes from the opposite signs in the definitions of the commutation relation. One immediately asks the question of why one introduces two polarizations (known as longitudinal) that shouldn't exist, one with peculiar creation/destruction operators. The answer is that if one calculates the probability for emitting into either of these unphysical states, the probabilities come out as equal and opposite. Thus, the negative probability is always canceled by additional positive probability, so that one can safely neglect the contribution from them if BOTH are left out of any calculation where they appear in the initial or final state. However, for intermediate states, such as above, they cannot be neglected. This particular form for the field operator is known as the "Feynmann Gauge". The simple fact that the Coulomb energy is always positive drives home the point that electromagnetic theory must incorporate some sort of peculiar behavior, whether that be in the form of having $\langle 0|aa^\dagger|0\rangle = -1$, or in introducing "ghost" fields (for another course).

$$\begin{aligned} 1. & \langle n, \vec{k} | H_{int} | 0 \rangle \\ &= \int d^3x_1 e^{i\vec{k} \cdot \vec{x}_1} \frac{1}{\sqrt{E_k}} \sqrt{\frac{2\pi\hbar^2 c^2}{V}} \rho(\vec{x}_1) \\ \delta E^{(2)} &= \sum_{\vec{k}} \frac{1}{E_k} \left| \int d^3x_1 e^{i\vec{k} \cdot \vec{x}_1} \frac{1}{\sqrt{E_k}} \sqrt{\frac{2\pi\hbar^2 c^2}{V}} \rho(\vec{x}_1) \right|^2 \\ &= \frac{2\pi\hbar^2 c^2}{V} \sum_{\vec{k}} \frac{1}{E_k^2} \int d^3x_1 d^3x_2 e^{i\vec{k} \cdot (\vec{x}_1 - \vec{x}_2)} \rho(\vec{x}_1) \rho(\vec{x}_2) \\ \sum_{\vec{k}} &\rightarrow \int \frac{V}{(2\pi)^3} d^3k, \quad E_k = \hbar c k \\ \delta E^{(2)} &= \left(\frac{1}{2\pi} \right)^2 \int \frac{d^3k}{k^2} \int d^3x_1 d^3x_2 \rho(\vec{x}_1) \rho(\vec{x}_2) e^{i\vec{k} \cdot (\vec{x}_1 - \vec{x}_2)} \\ 2. & \int \frac{d^3k}{k^2} e^{i\vec{k} \cdot (\vec{x}_1 - \vec{x}_2)} = 2\pi \int d\cos\theta dk e^{ik|\vec{x}_1 - \vec{x}_2|\cos\theta} \\ &= 4\pi \int \frac{dk}{k|\vec{x}_1 - \vec{x}_2|} \sin k|\vec{x}_1 - \vec{x}_2| = \frac{\pi}{|\vec{x}_1 - \vec{x}_2|} \int_{-\infty}^{\infty} \frac{du}{iu} (e^{iu} - e^{-iu}) \\ &= \frac{2\pi^2}{|\vec{x}_1 - \vec{x}_2|} \end{aligned}$$

$$\delta E^{(2)} = \frac{1}{2} \int d^3x_1 d^3x_2 \rho(\vec{x}_1) \rho(\vec{x}_2) \frac{1}{|\vec{x}_1 - \vec{x}_2|}$$