

Atomic Physics


PHY 215
Thermodynamics and
Modern Physics

Fall 2025
MSU

Outline: Atomic Physics in “1900”

- Classical Physics in “1900” (and now!)
- What the Chemists Knew
- Brownian Motion - atoms are real!
- Experiments:
 - New Particles: Electrons
 - New Radiation: X-rays & X-ray Diffraction
 - Elemental Transmutation: Radioactivity
- Atoms exist: how are they built?

Classical Physics




Sir Isaac Newton
1643-1727

Particles

$$\begin{aligned}\vec{F} &= m\vec{a} \\ \vec{F}_G &= -\frac{GMm}{r^2}\hat{r}\end{aligned}$$

(& Thermodynamics)



James Clerk Maxwell
1831-1879

$\vec{\nabla} \cdot \vec{E} = 4\pi\rho_e$
$\vec{\nabla} \cdot \vec{B} = 0$
$-\vec{\nabla} \times \vec{E} = \frac{\partial \vec{B}}{\partial t}$
$\vec{\nabla} \times \vec{B} = \frac{\partial \vec{E}}{\partial t} + 4\pi\vec{j}_e$

Waves


What the Chemists Knew

- Mass conserved in chemical reactions.
(Lavoisier)
- Law of Multiple Proportions: Masses of elements that combine, and react completely, form whole-number ratios.
(Dalton & Proust)
- Hydrogen is the lightest element.
 - Molecular weight: mass relative to H
- Atomic Hypothesis: everything made of discrete unites, atoms. **What is N_A ?**

Concept Test


- The Modern View: Atomic mass arises mostly from
 - Electrons
 - Protons
 - Neutrons
 - Protons and Neutrons, & $m_p \approx m_n$ ←

Brownian Motion




- Random motion of large particles suspended in a fluid medium.
- Einstein (1905): Average kinetic energy of *large particles* is $\frac{3}{2} kT$! (equipartition!)
- Can't measure K.E. directly. (freq. collisions!)
- Langevin (1909): $\langle x^2 \rangle = \frac{kT}{3\pi\eta a} t$ η =viscosity
 a =particle diameter
- Perrin (1909): measure, find k , deduce $N_A = R/k$.
Nobel Prize: 1926
- Establishes atomic hypothesis.

Experiments



Cathode Ray Tubes



Demo

Concept Test

- Cathode ray tubes used to be common elements of radios, televisions, and computers. Nowadays, these components have been replaced by
 - Resistors
 - Capacitors
 - Transistors ←
 - Power Supplies

The Electron!

J.J. Thomson (1856-1940)

- In 1897 he proved conclusively that cathode rays were indeed streams of **negatively-charged particles**.
- obtained better vacuum
- deflected the rays by both electric and magnetic fields.
- able to calculate

$$\frac{\text{Charge}}{\text{Mass}} = \frac{e}{m} = 1.7 \times 10^{11} \text{ Coulombs/kg}$$

(about 2000 times larger than that of Hydrogen atom)


large charge or
small mass!



J J Thomson
1856-1940
Nobel Prize: 1906

Image: <http://en.wikipedia.org>


Charge of the Electron



Oil Drop Experiment (1909)

Suspend charged oil drops falling in gravity with electrostatic field

- Measure Radius
- Suspend Drop in E field




Robert Millikan
1868-1953
Nobel Prize 1923

$$e = 1.602\,176\,53(14) \times 10^{-19} \text{ C}$$

electron charge: $-e$

X-Rays

- Roentgen (1895) studying cathode ray tube, noticed glass tubing glowing some distance away.
 - He covered the tube with black paper in a dark room. A fluorescent screen lit up, even if faced away from the discharge tube.
 - Then he placed his hand between the discharge tube and the screen and saw the shadow of his bones!
- ⇒ X-Rays! (later shown to be high frequency electromagnetic waves)



Wilhelm Röntgen

1845-1923

Nobel Prize 1901
(the first one!)




Images: <http://en.wikipedia.org>


X-Ray Scattering



Max von Laue
1879-1960
Nobel Prize 1914




© 2006 Brooks/Cole - Thomson



© 2006 Brooks/Cole - Thomson

$$\lambda_{\text{X-ray}} \approx \text{atomic spacing}$$

Bragg Diffraction



© 2006 Wadsworth - Thomson

William Lawrence Bragg


1890-1972

William Henry Bragg

1862-1942

Nobel Prize 1915

Crystals are a regular
array of atoms!



© 2006 Brooks/Cole - Thomson


Bragg's Law:

$$n\lambda = 2d \sin \theta$$

Images: Thornton and Rex

Example: DNA

- X-ray crystallography is used to determine the arrangement of atoms in a crystal.
- Sodium salt of DNA:



Franklin & Gosling,
Nature 171: 740-741



Rosalind Franklin
1920-1958



Francis Crick
1916-2004




James Watson
1928 -

Nobel Prize: 1962

Radioactivity I

- Becquerel (1896) was testing various fluorescent materials to see if they emitted X-rays.
- He sealed a photographic plate in black paper and sprinkled a layer of Uranyl Potassium sulfate onto the paper.
- He wanted to expose the salt to sunlight in order to make it fluoresce, but that day Paris was gray and overcast.
- Despite this images exposed with great intensity.
- The new phenomenon was named "radioactivity" by Marie Curie.




Antoine Henri Becquerel
1852-1908
Nobel Prize 1903

Radioactivity II

β -rays: electrons (Becquerel, 1900). They have a range of energies and are fast and penetrating. Can be absorbed by ~ 1 mm of lead.

α -rays: Helium nuclei (Rutherford, et al). They are heavy, slow, positively-charged particles. Absorbed by \sim few cm of air.

γ -rays: Electromagnetic radiation, with a higher frequency, lower wavelength, even than X-rays.



Transmutation

Pierre and Marie Curie found new radioactive elements, including Thorium, Polonium, and Radium.


A considerable amount of chemistry detective work, especially by the Curies, Rutherford, and Soddy led to a remarkable conclusion:

Every Radioactive decay is a transmutation of the elements, a change from one element to another.


Images: <http://en.wikipedia.org>
<http://www.ocrwm.doe.gov>



Marie Skłodowska-Curie
1867-1934
Nobel Prize 1903, 1911



Concept Test

- Elemental transmutation is induced on the MSU campus frequently at the following location:
 - Holmes Hall
 - NSCL  National Superconducting Cyclotron Laboratory
 - Biomedical Physical Sciences Building
 - Administration Building

Atoms Exist: Questions Remain!

- What is the structure of the atom?
 - Electrons and what?
 - Radioactivity and Elemental Transmutation?
- Do the laws of classical physics apply?

Understanding the answers to these questions
will take most of the rest of the semester!

Atomic Physics

PHY 215
Thermodynamics and
Modern Physics

Fall 2025
MSU

Outline

- Blackbody Radiation
- Planck's Formula
- Photoelectric Effect
 - X-Ray Production
- Compton Scattering
- Summary

Blackbody Radiation

As an object gets hot, it radiates energy.

How does the radiation depend on temperature?

What is the distribution among frequencies (or wavelengths)?

$\lambda \downarrow$
as $T \uparrow$

$I(\lambda, T)$ = Intensity as a function of wavelength λ and temperature T

Two experimental facts:

1) $\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m} \cdot \text{K}$

(Wien's displacement law)

2) Total Power (per unit Area)


$$R(T) = \int_0^{\infty} I(\lambda, T) d\lambda$$

given by

$$R(T) = \epsilon \sigma T^4$$

with $\epsilon=1$ for blackbody radiator
(Stefan-Boltzmann law)

$$\sigma = 5.67 \times 10^{-8} \text{ W} / (\text{m}^2 \cdot \text{K}^4)$$



Concept Test

- A blacksmith heats an iron rod which starts to radiate, beginning red, then orange, then yellow, and ultimately white. As the iron becomes hotter, the color changes because
 - Red light has a longer wavelength than blue light ←
 - Red and blue light have the same wavelength
 - Blue light has a longer wavelength than red light





Image: <http://www.saunalahti.fi>


Blackbody Spectrum

$I(\lambda, T)$



$$I(\lambda, T) = \frac{dE}{d\Omega dA d\lambda dt}$$

$$\neq \frac{2\pi ckT}{\lambda^4}$$



Lord Rayleigh and Sir James Jeans tried to explain the distribution using equipartition of energy: the average energy in each mode of oscillation of the radiation is $kT/2$.

Planck's Formula

Each oscillation mode can not absorb just any amount of energy. Each mode can only absorb energy in packets of fixed size.

It described the data perfectly!

The size of each energy packet (now termed **a quantum**) is proportional to the frequency of the mode:

$$E = h \nu$$

$$E \propto \text{Frequency!}$$

The proportionality factor (Planck's constant) is

$$h = 6.63 \times 10^{-34} \text{ J s.}$$

The above formula was first presented at a meeting of the German Physical Society on Dec. 14, 1900.

The birthday of Quantum Physics!

Light comes
in discrete
quanta! **Photons**



Max Planck

1848-1947

Nobel Prize 1918

$$I(\lambda, T) = \frac{2\pi c^2 h}{\lambda^5} \frac{1}{e^{hc/\lambda kT} - 1}$$

Planck's Constant

The energy in each frequency mode ν can only come in integer multiples of some fundamental unit (quanta):

$$E = h \nu$$

with

$$\begin{aligned} h &= 6.6261 \times 10^{-34} \text{ J} \cdot \text{s} \\ &= 4.1357 \times 10^{-15} \text{ eV} \cdot \text{s} \end{aligned}$$

$$1 \text{ V} = 1 \text{ J/C}$$

$$e \approx 1.6 \times 10^{-19} \text{ C}$$

$$1 \text{ eV} \approx 1.6 \times 10^{-19} \text{ J}$$

In a cathode ray tube, the change in potential energy of the electron is eV , where V is the voltage difference between cathode and anode.

Thus, eV is the Kinetic Energy of the electron when it hits the anode.


If all of this energy is converted into a single quantum of X-ray radiation (called a **photon**), then the photon would have a frequency given by

$$\nu = eV/h, \quad \lambda = c/\nu$$


This is the maximum frequency X-ray that could be emitted in the process.

For example, $2000 \text{ V} \rightarrow \lambda = 0.621 \text{ nm}$

Light Quanta?



Classically:
EM Radiation = Waves
Energy \propto (Amplitude)²



James Clerk Maxwell
1831-1879

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho_e$$

$$\vec{\nabla} \cdot \vec{B} = 0$$


$$-\vec{\nabla} \times \vec{E} = \frac{\partial \vec{B}}{\partial t}$$

$$\vec{\nabla} \times \vec{B} = \frac{\partial \vec{E}}{\partial t} + 4\pi\vec{j}_e$$

Photoelectric Effect I

1887 - Heinrich Hertz

Visible or UV light on metal surface may release electrons




Classical theory says energy of electrons should increase with intensity of light.

However, this was not the case.

Classically:

EM Radiation = Waves
Energy \propto (Amplitude)²



© 2006 Brooks/Cole - Thomson



Phillip Lenard
1862-1947
Nobel Prize 1905



Image: Thornton and Rex

Photoelectric Effect II


Experiments (Lenard) showed:

- 1) KE of photoelectrons depends only on ν (frequency) of light, independent of I (its intensity).
- 2) # of photoelectrons is proportional to I
- 3) For a given metal, there is a minimum ν , below which no photoelectrons are emitted.
- 4) The photoelectrons are emitted instantaneously, independent of I .

Classical theory could not explain these observations.



Concept Test

- In the photoelectric effect, if a fixed *frequency* of light shines on a metal, as the intensity of the light is increased
 - the number of electrons ejected increases 
 - the energy of the ejected electrons increases
 - both the number and energy of the ejected electrons increases

Photoelectric Effect III

1905 - Einstein explained:


Electromagnetic radiation transferred in discrete bundles of energy ("photons").
The energy of each photon is

$$E = h \nu$$

The KE of an emitted photoelectron is

$$KE = h \nu - \phi$$

- $h\nu$ is Energy of the incident photon
- ϕ is the binding energy of electron to the metal surface (the work function).



Albert Einstein


1879-1955

Nobel Prize 1921

Einstein extended Planck's hypothesis
to explain the photoelectric effect!

X-Ray Production

Inverse photoelectric effect:




© 2006 Brooks/Cole - Thomson

$$eV_0 = hf_{\max} = \frac{hc}{\lambda_{\min}}$$

Duane-Hunt limit

$$\lambda_{\min} = \frac{hc}{eV_0} = \frac{1.240 \times 10^{-6} \text{ V} \cdot \text{m}}{V_0}$$

Maximum frequency
related to maximum energy ✓




© 2006 Brooks/Cole - Thomson

Images: Thornton and Rex

Compton Scattering I

Classical Picture of Scattering of Light:


Wave picture:



Electron accelerates,
and reradiates

Light scatters off all electrons in metal, and
scattered frequency same as incident frequency


Concept Test



- Light is a transverse wave: the direction of the oscillating E and B fields is perpendicular to the direction of propagation. What is wrong with the diagram above?
- Light doesn't have to be blue
- The electron is moving in the wrong direction ←
- The scattered light is not a spherical wave
- Electrons aren't green


Compton Scattering II

Quantum picture:



A quantum of light scatters off an electron, and can change energy and hence frequency


Compton Formula



$$\lambda' - \lambda = \frac{h}{m_e c} (1 - \cos \theta)$$

$$\frac{1}{E_\gamma} - \frac{1}{E'_\gamma} = \frac{1}{m_e c^2} (1 - \cos \theta)$$

(You will derive this in homework.)



Arthur Holly Compton
1892-1962
Nobel Prize 1927

Summary

- Planck's formula, $E=h\nu$, explains
 - Blackbody Spectrum
 - Photoelectric Effect
 - X-ray Production
 - Compton Scattering

Light behaves both like a particle and a wave!

Light “particle” - Photon

Atomic Physics

PHY 215
Thermodynamics and
Modern Physics

Fall 2025
MSU


Outline

- Waves vs. Particles
 - Review of Wave Diffraction
- deBroglie Waves
- Davison and Germer
- Electron Double Slit
- Electrons Behave Like Waves!
- Complementarity



Particles vs. Waves

Property	Particle	Wave
Location	Definite	Indefinite
Momentum	Definite	Indefinite
Interference	No	Yes

Particle




Wave



Diffraction, Thomas Young (1803)

Images: <http://en.wikipedia.org>
<http://www.staff.amu.edu.pl>
<http://micro.magnet.fsu.edu>

Bragg Diffraction



© 2006 Wadsworth - Thomson


William Lawrence Bragg

1890-1972

William Henry Bragg

1862-1942

Nobel Prize 1915



© 2006 Brooks/Cole - Thomson

Bragg's Law:


$$n\lambda = 2d \sin \theta$$

Classically, light
behaves as a wave!

Images: Thornton and Rex

Electrons


- Are parts of an atom.
- Atoms are particles. —————→
- Ergo, electrons are particles.
- Or are they...



Graphite

Concept Test

- For a light wave with wavelength λ and frequency ν , the following relation is always true (in a vacuum):

- $\lambda \nu = c$  $[c] = \frac{\text{m}}{\text{s}} = [\lambda] \cdot [\nu]$


- $\lambda c = \nu$

- $\nu c = \lambda$

where c is the speed of light.

For Light

- $\lambda\nu=c$
- $E=pc$ (Einstein)
- $E=h\nu$ (Planck)
- $pc=h\nu=hc/\lambda$
- $\therefore \lambda=h/p$



A Crazy Idea...

Matter waves

An electron with momentum p
“has” a wavelength

$$\lambda = h/p \quad !$$

Quantum “fuzziness”: a wave is hard to
localize within a size of order λ


Heisenberg will make this more precise!




Louis de Broglie
1892-1987
Nobel Prize: 1929

Experimental Proof!

1925: Diffraction of Electrons by a Nickel Crystal



Intensity = radial distance along dashed line to data at angle ϕ



Clinton Davission (R)

(1881-1958)


Nobel Prize 1937

Lester Germer

(1896-1971)

Images: Thornton and Rex


What is the wavelength of a tennis ball?


$$\begin{aligned} \lambda &= \frac{6.63 \times 10^{-34} \text{ J s}}{60 \frac{\text{mi}}{\text{hr}} \cdot 1600 \frac{\text{m}}{\text{mi}} \cdot \frac{1}{3600} \frac{\text{hr}}{\text{s}} \cdot 0.1 \text{ kg}} \\ &= 2.5 \times 10^{-34} \text{ m} \\ &= \mathcal{O}(10^{-24}) \times \text{size of atom} \end{aligned}$$

Quantum “fuzziness” of tennis ball is *very* small!



(I can’t blame my inability to hit a tennis ball to quantum effects.)

Concept Test

- The de Broglie wavelength of a tennis ball, $\lambda = h/p$, goes to ∞ (the “fuzziness” of the ball) as $p \rightarrow 0$. Why doesn't this prevent us from picking up tennis balls left on the court?
- de Broglie's formula doesn't apply
- $h \rightarrow 0$ when the ball isn't moving
- p is never exactly 0 


Double Slit Experiment

Light



Diffraction, Thomas Young (1803)

Double Slit Experiment: Electrons!



© 2006 Brooks/Cole - Thomson

C. Jönsson, 1961





Image: Thornon and Rex
<http://stephenwhitt.files.wordpress.com>

One electron at a time!



Electrons, even one at a time, behave like waves!



Hitachi Laboratories

Dr.Tonomura Akiro

Movie: www.hitachi.com

Duality and Complementarity

- Wave-Particle Duality: all matter and energy exhibits *both* wave and particle like properties!
- Complementarity: a single quantum-mechanical system can behave like a wave or a particle, *but not both simultaneously!* (Bohr)

Summary, so far

- Light is classically a wave, but can behave like a particle (e.g. the photoelectric effect).
- Electrons are classically described as particles, but can behave like waves (e.g. the Davisson-Germer experiment).
- We need a description that unifies the particle and wave aspects of natural systems!