
nagytimo@msu

Nagy,

Tibor

Keep this exam **CLOSED** until advised by the instructor.

120 minute long closed book exam.

Fill out the bubble sheet: last name, first initial, **student number (PID)**. Leave the section, code, form and signature areas empty.

Four two-sided handwritten 8.5 by 11 help sheets are allowed.

When done, hand in your **test** and your **bubble sheet**.

Thank you and good luck!

Possibly useful constants:

- $g = 9.81 \text{ m/s}^2$
- $G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$
- $\rho_{\text{water}} = 1000 \text{ kg/m}^3 = 1 \text{ kg/l} = 1 \text{ g/cm}^3$
- $1 \text{ atm} = 101.3 \text{ kPa} = 760 \text{ mmHg}$
- $N_A = 6.02 \times 10^{23} \text{ 1/mol}$
- $R = 8.31 \text{ J/(molK)}$
- $k_B = 1.38 \times 10^{-23} \text{ J/K}$
- $c_{\text{water}} = 4.1868 \text{ kJ/(kg}^\circ\text{C)} = 1 \text{ kcal/(kg}^\circ\text{C)}$
- $1 \text{ cal} = 4.1868 \text{ J}$
- $\sigma = 5.67 \times 10^{-8} \text{ W/(m}^2\text{K}^4)$
- $b = 2.90 \times 10^{-3} \text{ m}\cdot\text{K}$

Possibly useful Moments of Inertia:

- Solid homogeneous cylinder: $I_{\text{CM}} = (1/2)MR^2$
 - Solid homogeneous sphere: $I_{\text{CM}} = (2/5)MR^2$
 - Thin spherical shell: $I_{\text{CM}} = (2/3)MR^2$
 - Straight thin rod with axis through center: $I_{\text{CM}} = (1/12)ML^2$
 - Straight thin rod with axis through end: $I = (1/3)ML^2$
-

Please, sit in seat:

Thank you!


1 pt Are you sitting in the seat assigned?

1.A Yes, I am.

$\frac{4}{4}$ pt A large rock is released from rest from the top of a tall building. The average speed of the rock during the first second of the fall is 5 m/s. What is the average speed of the rock during the third second? (In this question we use the approximate value of 10 m/s² for the gravitational acceleration.)

2. **A** 0 m/s
B 5 m/s
C 25 m/s
D 30 m/s
E 100 m/s
F 15 m/s
G 50 m/s
H 20 m/s
I 10 m/s

A car is waiting at an intersection. When the traffic light turns green, the car starts moving. After some time the car comes to rest at another traffic light. The figure below shows the velocity of the car as a function of time.



One can clearly identify three different stages of this motion.

$\frac{2}{2}$ pt What is the acceleration of the car during the second stage of the motion?
(in m/s²)

3. **A** -2.00 **B** -1.00 **C** -0.667 **D** 0
E 0.400 **F** 0.500 **G** 0.667 **H** 2.00

$\frac{2}{2}$ pt What is the total distance travelled by the car between the two traffic lights?
(in m)

4. **A** 26.4 **B** 35.1 **C** 46.6 **D** 62.0
E 82.5 **F** 110. **G** 146 **H** 194

A baseball is projected horizontally with an initial speed of 17.9 m/s from a height of 2.33 m. (Please, neglect air friction in this question.)


2 pt What is the horizontal component of the ball's velocity, when the ball hits the ground?
(in m/s)

5. **A** 1.53×10^1 **B** 1.79×10^1 **C** 2.09×10^1
 D 2.45×10^1 **E** 2.87×10^1 **F** 3.35×10^1
 G 3.92×10^1 **H** 4.59×10^1

2 pt What is the magnitude of the vertical component of the ball's velocity, when the ball hits the ground?
(in m/s)

6. **A** 3.22 **B** 4.66 **C** 6.76
 D 9.80 **E** 1.42×10^1 **F** 2.06×10^1
 G 2.99×10^1 **H** 4.33×10^1

Two masses, $m_1 = 3.20$ kg and $m_2 = 5.98$ kg are on a horizontal frictionless surface and they are connected together with a rope as shown in the figure.




2 pt The rope will snap if the tension in it exceeds 75.0 N. What is the maximum value of the force F which can be applied?
(in N)

7. **A** 3.89×10^1 **B** 5.17×10^1 **C** 6.88×10^1
 D 9.15×10^1 **E** 1.22×10^2 **F** 1.62×10^2
 G 2.15×10^2 **H** 2.86×10^2

2 pt What is the acceleration of the whole system, when this maximum force is applied?
(in m/s^2)

8. **A** 2.34×10^1 **B** 2.65×10^1 **C** 2.99×10^1
 D 3.38×10^1 **E** 3.82×10^1 **F** 4.32×10^1
 G 4.88×10^1 **H** 5.51×10^1

3 pt A block is at rest on a frictional incline. (See figure.)



Which vector best represents the direction of the force exerted by the surface on the block?


9. A
 B
 C
 D
 E
 F
 G
 H
 I: the force is zero.

4 pt An athlete, swimming at a constant speed, covers a distance of 226 m in a time period of 5 minutes and 31 seconds. The drag force exerted by the water on the swimmer is 70.0 N. What is the power the swimmer must provide to overcome that drag force.

(in W)

10. A 1.57×10^1 B 2.27×10^1 C 3.30×10^1
 D 4.78×10^1 E 6.93×10^1 F 1.00×10^2
 G 1.46×10^2 H 2.11×10^2

On a roller coaster ride the total mass of a cart - with two passengers included - is 316 kg. Peak **K** is at 46.6 m above the ground and peak **L** is at 24.0 m. At location **K** the speed of the cart is 15.3 m/s, and at location **L** it is 12.4 m/s. (The wheel mechanism on roller coaster carts always keeps the carts safely on the rail.)



4 pt How much mechanical energy is lost due to friction between the two peaks?

(in J)


11. A 8.90×10^3 B 1.29×10^4 C 1.87×10^4
 D 2.71×10^4 E 3.94×10^4 F 5.71×10^4
 G 8.28×10^4 H 1.20×10^5

4 pt A 927 kg automobile slides across an icy street at a speed of 62.7 km/h and collides with a parked car. The two cars lock up and they slide together with a speed of 29.7 km/h. What is the mass of the parked car?

(in kg)

12. A 6.59×10^2 B 8.24×10^2 C 1.03×10^3
 D 1.29×10^3 E 1.61×10^3 F 2.01×10^3
 G 2.51×10^3 H 3.14×10^3

2 pt A mass of $m = 1.41$ kg connected to a spring oscillates on a horizontal frictionless surface as shown in the figure.



The equation of motion of the mass is given by

$$x = 0.333 \cos(1.02t)$$

where the position x is measured in meters, the time t in seconds. Determine the period of the motion.

(in s)

13. A 5.45 B 6.16 C 6.96
 D 7.87 E 8.89 F 1.00×10^1
 G 1.13×10^1 H 1.28×10^1

2 pt What is the maximum speed reached by the mass?

(in m/s)


14. A 1.84×10^{-1} B 2.08×10^{-1} C 2.35×10^{-1}
 D 2.66×10^{-1} E 3.01×10^{-1} F 3.40×10^{-1}
 G 3.84×10^{-1} H 4.34×10^{-1}

2 pt Determine the spring constant.

(in N/m)


15. A 4.81×10^{-1} B 6.01×10^{-1} C 7.51×10^{-1}
 D 9.39×10^{-1} E 1.17 F 1.47
 G 1.83 H 2.29

4 pt An extended body (not shown in the figure) has its center of mass (CM) at the origin of the reference frame. In the case below give the direction for the torque τ with respect to the CM on the body due to force \mathbf{F} acting on the body at a location indicated by the vector \mathbf{r} .



16. A X B -X C Y D -Y E Z
 F -Z

10 pt A small mass M attached to a string slides in a circle (Y) on a frictionless horizontal table, with the force \mathbf{F} providing the necessary tension (see figure). The force is then decreased slowly and then maintained constant when M travels around in circle (X). The radius of circle (X) is twice the radius of circle (Y).



▷ As M moves from Y to X, the work done by \mathbf{F} is 0.

17. A true B false C greater than
 D less than E equal to

▷ While going from Y to X, there is a torque on M .

18. A true B false C greater than
 D less than E equal to

▷ M 's kinetic energy at X is half that at Y.

19. A true B false C greater than
 D less than E equal to


▷ M 's angular momentum at X is that at Y.

20. A true B false C greater than
 D less than E equal to

▷ M 's angular velocity at X is half that at Y.

21. A true B false C greater than
 D less than E equal to

A crate with a mass of $M = 84.5$ kg is suspended by a rope from the endpoint of a uniform boom. The boom has a mass of $m = 135$ kg and a length of $l = 7.25$ m. The midpoint of the boom is supported by another rope which is horizontal and is attached to the wall as shown in the figure.



2 pt The boom makes an angle of $\theta = 65.0^\circ$ with the vertical wall. Calculate the tension in the vertical rope.
(in N)

22. A 6.23×10^2 B 8.29×10^2 C 1.10×10^3
 D 1.47×10^3 E 1.95×10^3 F 2.59×10^3
 G 3.45×10^3 H 4.59×10^3

2 pt What is the tension in the horizontal rope?
(in N)

23. A 5.12×10^3 B 6.40×10^3 C 7.99×10^3
 D 9.99×10^3 E 1.25×10^4 F 1.56×10^4
 G 1.95×10^4 H 2.44×10^4

4 pt A 230 kg satellite is orbiting Earth on a circular orbit with a speed of 5.71 km/s. Determine the height of the satellite above Earth's surface. (The mass of the Earth is 5.97×10^{24} kg, and the radius of the Earth is 6370 km.)
(in km)

- 24.** A 3.30×10^3 B 4.39×10^3 C 5.84×10^3
D 7.77×10^3 E 1.03×10^4 F 1.37×10^4
G 1.83×10^4 H 2.43×10^4
-

4 pt Glucose solution is administered to a patient in a hospital. The density of the solution is 1.300 kg/l. If the blood pressure in the vein is 25.7 mmHg, then what is the minimum necessary height of the IV bag above the position of the needle?
(in cm)

- 25.** A 12.8 B 18.5 C 26.9 D 38.9
E 56.5 F 81.9 G 118.7 H 172.2
-

4 pt An object weighs 95.9 N in air. When it is suspended from a force scale and completely immersed in water the scale reads 24.4 N. Determine the density of the object.
(in kg/m^3)

- 26.** A 4.40×10^2 B 5.49×10^2 C 6.87×10^2
D 8.58×10^2 E 1.07×10^3 F 1.34×10^3
G 1.68×10^3 H 2.10×10^3
-

4 pt An Airbus A380-800 passenger airplane is cruising at constant altitude on a straight line with a constant speed. The total surface area of the two wings is 395 m^2 . The average speed of the air just below the wings is 252 m/s, and it is 278 m/s just above the surface of the wings. What is the mass of the airplane? The average density of the air around the airplane is $\rho_{\text{air}} = 1.11 \text{ kg/m}^3$.
(in kg)

- 27.** A 2.725×10^5 B 3.079×10^5
C 3.480×10^5 D 3.932×10^5
E 4.443×10^5 F 5.021×10^5
G 5.674×10^5 H 6.411×10^5
-

4 pt Two sounds have intensities of 1.80×10^{-8} and 6.90×10^{-4} W/m² respectively. What is the magnitude of the sound level difference between them in dB units?

28. A 6.23 B 8.28 C 11.01 D 14.65
E 19.48 F 25.91 G 34.46 H 45.84
-

4 pt A bag filled with lead shots is dropped from a height of $h = 25.1$ m. The total mass of the bag is $m = 515$ g. What is the increase in the temperature of the lead shots, after the bag hits the ground? (The specific heat of lead is $c = 130$ J/kgK.)
(in K)

29. A 7.76×10^{-1} B 9.70×10^{-1} C 1.21
D 1.52 E 1.89 F 2.37
G 2.96 H 3.70
-

4 pt What is the pressure of 2.03 moles of Nitrogen gas in a 7.85 liter container, if the temperature of the gas is 30.5 °C?
(in atm)

30. A 3.06 B 4.44 C 6.44 D 9.34
E 13.54 F 19.64 G 28.47 H 41.29
-

10 pt Constant amount of ideal gas is kept inside a cylinder by a piston. Then the gas is compressed **isobarically**. Compare the initial (i) and the final (f) physical quantities of the gas to each other.

▷ The pressure p_f is ... p_i .
31. A equal to B less than
 C greater than

▷ The internal energy U_f is ... U_i .
32. A equal to B less than
 C greater than

▷ The temperature T_f is ... T_i .
33. A equal to B less than
 C greater than

▷ The volume V_f is ... V_i .
34. A equal to B less than
 C greater than

▷ The entropy S_f is ... S_i .
35. A equal to B less than
 C greater than

3 pt A Stirling-engine is used in the heat-pump mode to heat a house. The engine maintains a temperature of 23.7 °C inside the house. The temperature of the Earth loop is 11.3 °C. (The Earth loop buried deep under the ground is the cold reservoir of this heat pump.) What is the coefficient of performance of this heat pump?

- 36.** A 6.28 B 7.84 C 9.81
 D 1.23×10^1 E 1.53×10^1 F 1.92×10^1
 G 2.39×10^1 H 2.99×10^1
-

3 pt If the power of the electric motor driving the heat pump is 174 W, then what is the rate at which heat is delivered to the house?

(in W)

- 37.** A 4.48×10^2 B 6.50×10^2 C 9.42×10^2
 D 1.37×10^3 E 1.98×10^3 F 2.87×10^3
 G 4.17×10^3 H 6.04×10^3
-